

Hyperbolic manifolds with polyhedral boundary

Jean-Marc Schlenker*

Nov. 2001; revised, Sept. 2002

Abstract

Let $(M, \partial M)$ be a compact 3-manifold with boundary which admits a complete, convex co-compact hyperbolic metric. For each hyperbolic metric g on M such that ∂M is smooth and strictly convex, the induced metric on ∂M has curvature $K > -1$, and each such metric on ∂M is obtained for a unique choice of g . A dual statement is that, for each g as above, the third fundamental form of ∂M has curvature $K < 1$, and its closed geodesics which are contractible in M have length $L > 2\pi$. Conversely, any such metric on ∂M is obtained for a unique choice of g .

We are interested here in the similar situation where ∂M is not smooth, but rather looks locally like an ideal polyhedron in H^3 . We can give a fairly complete answer to the question on the third fundamental form — which in this case concerns the dihedral angles — and some partial results about the induced metric.

This has some by-products, like an affine piecewise flat structure on the Teichmüller space of a surface with some marked points, or an extension of the Koebe circle packing theorem to many 3-manifolds with boundary.

Résumé

Soit $(M, \partial M)$ une variété compacte de dimension 3 à bord, qui admet une métrique complète convexe co-compacte. Pour chaque métrique hyperbolique g sur M telle que ∂M est régulier et strictement convexe, la métrique induite sur ∂M est à courbure $K > -1$; réciproquement, chaque métrique à courbure $K > -1$ sur ∂M est obtenue pour un unique choix de g . Un énoncé dual est que, pour ces métriques g sur M , la troisième forme fondamentale de ∂M est à courbure $K < 1$, et ses géodésiques fermées qui sont contractiles dans M sont de longueur $L > 2\pi$; réciproquement, chaque métrique de ce type est obtenue pour un unique choix de g .

Nous nous intéressons au cas similaire où ∂M n'est pas régulière, mais ressemble au contraire localement à un polyèdre idéal dans H^3 . On donne un énoncé assez complet concernant la troisième forme fondamentale — qui dans ce cas se formule en termes d'angles dièdres — et un énoncé partiel pour la métrique induite sur le bord.

Ceci a comme conséquence l'existence d'une métrique affine plate par morceaux sur l'espace de Teichmüller d'une surface munie de points marqués, ou une extension du théorème de Koebe sur les empilements de cercles à beaucoup de 3-variétés à bord.

Contents

1	Introduction	3
1.1	Hyperbolic manifolds with convex boundary	3
1.2	Circle packings	5
1.3	Induced metrics on the boundary	5
1.4	Fuchsian polyhedra	5
1.5	Some hints on the techniques	6
1.6	What follows	7

*Laboratoire Emile Picard, UMR CNRS 5580, Université Paul Sabatier, 118 route de Narbonne, 31062 Toulouse Cedex 4, France. schlenker@picard.ups-tlse.fr; <http://picard.ups-tlse.fr/~schlenker>.

2	The Schläfli formula	7
2.1	A symplectic viewpoint	8
2.2	Ideal polyhedra	8
2.3	The volume function	9
3	Convex hulls of ideal points	10
3.1	Some definitions	10
3.2	Relation with circle packings	10
3.3	Existence of the bent case	11
3.4	Necessary conditions on convex surfaces	12
3.5	Third fundamental form versus dihedral angles	14
4	The fuchsian case	15
4.1	Infinitesimal rigidity of finite fuchsian polyhedra	16
4.2	Compactness of fuchsian polyhedra	17
4.3	Induced metrics on finite polyhedra	22
4.4	Fuchsian ideal manifolds	23
5	Triangulations	24
5.1	Triangulations, cellulations	24
5.2	From a cellulation to a triangulation	26
5.3	Some elementary combinatorics	28
6	Hyperbolic structures on triangulated manifolds	28
6.1	Sheared hyperbolic structures	29
6.2	Angles	29
7	First order variation of the volume	30
7.1	Definitions and first properties	30
7.2	Volume differential and shears	31
7.3	Consequences	33
8	Dihedral angles	34
8.1	Compactness of ideal manifolds	34
8.2	Dihedral angles with a given triangulation	36
8.3	Fixed triangulations of the boundary	37
8.4	Spaces of angle assignations	37
8.5	Unique realization for some dihedral angle assignations	38
8.6	Proof of the main theorem	39
8.7	An affine structure on some Teichmüller spaces	40
8.8	Hyperbolic polyhedra	42
9	Induced metrics	42
9.1	Finite area metrics on surfaces	42
9.2	The lengths of the edges of σ	43
9.3	Rigidity and the Schläfli formula	44
9.4	The fuchsian case	44
9.5	Ideal polyhedra in H^3	45
10	Circles packings	45
10.1	From dihedral angles to circle configurations	45
10.2	Circles packings and CP^1 -structures	47

11 Concluding remarks and questions	47
11.1 Manifolds with cusps	47
11.2 Hyperideal polyhedra	47
11.3 Complete manifolds of finite volume	48
11.4 Convex cores	48
11.5 Orbifolds, cone-manifolds	49
11.6 Affine structures	49
11.7 Smooth boundaries	49
11.8 Higher dimensions	49

1 Introduction

1.1 Hyperbolic manifolds with convex boundary

We will be motivated by the following results from [Sch02a], which show that the hyperbolic metrics on M such that ∂M is smooth and strictly convex can be well understood from quantities induced on the boundary.

Theorem 1.1. *Let g be a hyperbolic metric on M such that ∂M is smooth and strictly convex. Then the induced metric I on ∂M has curvature $K > -1$. Each smooth metric on ∂M with $K > -1$ is induced on ∂M for a unique choice of g .*

Theorem 1.2. *Let g be a hyperbolic metric on M such that ∂M is smooth and strictly convex. Then the third fundamental form III of ∂M has curvature $K < 1$, and its closed geodesics which are contractible in M have length $L > 2\pi$. Each such metric is obtained for a unique choice of g .*

Theorem 1.1 was conjectured by Thurston, and its existence part was proved by Labourie [Lab92]. Note that the hypothesis of theorem 1.2 is that the metric on ∂M , when lifted to the boundary of the universal cover of M , is globally CAT(1); this implies a local "curvature" condition, but also a global condition, namely that all closed geodesics have length larger than 2π .

A striking feature of both those statements is that they appear to hold, at least in some cases, without any smoothness assumption (beyond the convexity). In the simple case where M is a ball, both results can be stated in terms of convex surfaces in H^3 :

Theorem 1.3 (Pogorelov [Pog73]). *Let h be a smooth metric with curvature $K > -1$ on S^2 . Then (S^2, h) has a unique isometric embedding into H^3 .*

Theorem 1.4 ([Sch96, Sch94]). *Let h be a smooth metric with curvature $K < 1$ on S^2 . Then (S^2, h) has an embedding in H^3 such that the third fundamental form of the image is h if and only if all closed geodesics of (S^2, h) have length $L > 2\pi$. This embedding is then unique.*

Both of those statements also hold without any smoothness assumption; it is a result of [Pog73] for theorem 1.3, and has been announced recently by G. Moussong for theorem 1.4 (see [DM99]). In particular, the polyhedral case correspond to earlier results of Aleksandrov [Ale58] for theorem 1.3, and to results of Andreev [And70], Rivin and Hodgson [Riv86, RH93] for theorem 1.4.

It also appears that theorems 1.1 and 1.2 hold, at least in some cases, for convex surfaces which are not complete. This is again most apparent when M is a ball. In particular:

- direct analogs of theorems 1.1 and 1.2 hold for smooth, non-compact convex surfaces in H^3 [Sch98b].
- the analog of theorem 1.1 is true for ideal polyhedra, a result of Rivin [Riv92].
- the analog of theorem 1.2 also holds for ideal polyhedra, a result of Andreev [And71] and Rivin [Riv96]. Thurston [Thu97] noted that those results are also strongly related to questions on circle packings, in particular the Koebe theorem.
- the analog of theorem 1.1 holds for hyperideal polyhedra in H^3 , see [Sch98a].

- theorem 1.2 also holds for hyperideal polyhedra, see [Sch98a, BB02]

When M is topologically more complicated than a ball, known results are limited to the "fuchsian" case, i.e. when M is the product of a surface of genus at least 2 by an interval, with an isometric involution. In this case, a version of theorem 1.2, concerning fuchsian manifolds whose boundary is locally like an ideal polyhedron, can be found hidden behind work of Thurston [Thu97, chapter 13] and Colin de Verdière [CdV91] on circle packings on hyperbolic surfaces; the relationship (which is well known) should be clear from section 3 below.

The main goal of this paper is to extend this result to the situation where M is topologically "general" — but its boundary will always be supposed to be locally like an ideal polyhedron, in a sense which is precisely defined in section 3. In this case, we will call M an **ideal hyperbolic manifold**.

We need the following definition. A **cellulation** of ∂M is a decomposition of ∂M into the images by diffeomorphisms of convex polygons in $\mathbf{R}^2$. We will call the images of the polygons the **2-cells** of the cellulations, and the **1-cells** will be the images of the edges of the polygons. We demand that the cellulations are well behaved in the sense that:

- the 2-cells have disjoint interiors.
- two 1-cells which are distinct have disjoint interiors.
- the intersection of two 2-cells is a disjoint union of 1-cells and vertices.
- at least 3 images of polygons meet at the image of each vertex.

Definition 1.5. Let Γ be the 1-skeleton of a cellulation of ∂M . A **circuit** in Γ is a sequence $e_0, e_1, \dots, e_n = e_0$ of edges of Γ such that the dual edges $e_0^*, e_1^*, \dots, e_n^*$ are the successive edges of a closed path in the dual graph Γ^* , which is contractible in M . A circuit is **elementary** if the dual path bounds a face (i.e. containing no edge or vertex).

We can now describe the dihedral angles of ideal hyperbolic manifolds; it extends the results of [And71, Riv96] on ideal polyhedra.

Theorem 8.15. Suppose that M has incompressible boundary. Let Γ be the 1-skeleton of a cellulation of ∂M . Let w be a function from the edges of Γ to $(0, \pi)$ such that:

1. for each elementary circuit in Γ , the sum of the values of w is 2π ;
2. for each non-elementary circuit in Γ , the sum of the values of w is strictly larger than 2π .

Then there is a unique hyperbolic metric g on M , such that (M, g) is an ideal hyperbolic manifold, with exterior dihedral angles given by w .

The hypothesis that M has incompressible boundary is necessary for technical reasons (see section 8). Theorem 1.1 would seem to indicate that the result also holds without this hypothesis.

Note that the hypothesis of this theorem on the function w , implies topological properties on Γ . In particular:

Lemma 1.6. Let σ be a cellulation of ∂M , such that there is a function w on the edges of σ , satisfying the hypothesis of theorem 8.15. Then:

- each 1-cell, when lifted to the universal cover of ∂M , is topologically a segment.
- each 2-cell, when lifted to the universal cover of ∂M , is topologically a disk.

The proof is in section 3.

1.2 Circle packings

Theorem 8.15 has a consequence in terms of circle packing on the boundary of 3-manifolds. Remember that a **circle packing** on the sphere S^2 is a set of closed disks with disjoint interiors in S^2 . Given a circle packing, one can define its **incidence graph** as a graph on S^2 which has one vertex for each disk, and an edge between two vertices if and only if the corresponding disk are tangent.

The classical Koebe circle packing theorem states that, for each graph γ in S^2 which is the 1-skeleton of a triangulation, there exists a unique circle packing whose incidence graph is Γ . It was proved by Koebe [Koe36] for triangulations, and extended by Thurston [Thu97] using the Andreev theorem on ideal polyhedra [And71]; in this more general case one should demand that, for any connected component of the complement of the disks, there is a circle which is orthogonal to all the neighboring disks.

Thurston also extended this theorem to circle packings on hyperbolic surfaces. Theorem 8.15 provides a further extension to the boundary of our 3-manifold M , when it is provided with the CP^1 -structure on the boundary coming from a hyperbolic metric on M .

Theorem 10.2. *Suppose that M has incompressible boundary. Let Γ be the 1-skeleton of a triangulation of ∂M . There is a unique couple (g, c) , where g is a complete, convex co-compact hyperbolic metric on M , and c is a circle packing on ∂M (for the CP^1 -structure defined on ∂M by g) whose incidence graph is Γ .*

If one considers not only triangulation but more generally cellulations, the same result holds, with the additional condition that, for each connected component of the complement of the closed disks, there exists a circle which is orthogonal to all the adjacent circles — this condition is automatically satisfied for a triangulation.

The proof is given in section 10.

1.3 Induced metrics on the boundary

It would be interesting to know whether theorem 1.1 extends to ideal hyperbolic manifolds. Unfortunately I do not know the answer to this question; but an infinitesimal version does hold:

Lemma 1.7. *For each ideal hyperbolic manifold M , each infinitesimal variation of the hyperbolic structure on M (among ideal hyperbolic manifolds) induces a non-trivial infinitesimal variation of the induced metric on ∂M .*

This assertion is worth mentioning because, in other questions related to theorem 1.1, it is precisely this infinitesimal rigidity which is lacking (see section 1).

However, in this case it is not sufficient to obtain a global result concerning the induced metrics on the boundary of ideal hyperbolic manifolds, like in theorem 8.15 for the third fundamental form. The reason is that, to understand the induced metrics completely, one would have to go beyond the category of ideal hyperbolic manifolds, and obtain an infinitesimal rigidity result also for "bent" manifolds (as defined in section 3). We give more details on this in section 11.

We will also give below a more precise result in the special case of manifolds which we call "fuchsian"; they are ideal hyperbolic manifolds such that ∂M has two connected component, with an isometric involution exchanging the connected components of ∂M .

1.4 Fuchsian polyhedra

In the course of the proof of this theorem, we are led to study fuchsian equivariant polyhedra; they are the objects arising as the universal covers of the boundaries of the fuchsian manifolds mentioned above. As a consequence of theorem 8.15, we find a characterization of the dihedral angles of fuchsian ideal polyhedra. In addition, we will give in section 4 results concerning other fuchsian polyhedra, having some non-ideal vertices.

Definition 1.8. *A fuchsian polyhedron is a triple (S, ϕ, ρ) , where:*

- S is a surface of genus $g \geq 2$, with N marked points $x_1, x_2, \dots, x_N$, $N \geq 1$.

- ϕ is a polyhedral map from the universal cover $\tilde{S}$ of S into $H^3 \cup \partial_\infty H^3$ such that the image is locally like a polyhedron in H^3 , with vertices the images by ϕ of the inverse images in $\tilde{S}$ of the x_i .
- ρ is a morphism from $\pi_1 S$ into the subgroup of the isometry group of H^3 of isometries fixing a totally geodesic 2-plane P_0 .
- for any $x \in \tilde{S}$ and any $\gamma \in \pi_1 S$, $\phi(\gamma x) = \rho(\gamma)\phi(x)$.

Some special cases are of interest. Let I be the set of points in $\tilde{S}$ which are sent by the canonical projection $\tilde{S} \rightarrow S$ to a marked point. We say that (S, ϕ, ρ) is:

- ideal, if each point in I is sent to an ideal point.
- finite, if no point in I is sent to an ideal point.
- semi-ideal, if the points in I can be sent to either ideal or non-ideal vertices (this includes the finite and the ideal cases).

We obtain an existence result for the third fundamental forms of fuchsian polyhedra.

Theorem 4.25. *Let Σ be a surface of genus $g \geq 2$, and let h be a spherical cone-metric on Σ , with negative singular curvature at the singular points. Suppose that all contractible closed geodesics of (Σ, h) have length $L > 2\pi$, except when they bound a hemisphere. Then there is a unique fuchsian polyhedral embedding of (Σ, h) into H^3 whose third fundamental form is h .*

As a consequence of the analysis of the dihedral angles of ideal hyperbolic manifolds in the fuchsian case, we will obtain the following.

Theorem 8.19. *For each $g \geq 2$ and each $N \geq 1$, there is a natural unimodular piecewise affine structure $\mathcal{A}_{g,N}$ on the Teichmüller $\mathcal{T}_{g,N}$ space of the genus g surface with N marked points.*

This affine structure has some singularities, but also a number of interesting properties; for instance there is a natural function on $\mathcal{T}_{g,N}$, which is defined by taking the volume of a fuchsian ideal hyperbolic manifold, and it is concave on the maximal dimension cells of $\mathcal{A}_{g,N}$. There are also some questions which remain open concerning $\mathcal{A}_{g,N}$.

Finally, in the case of fuchsian polyhedra, the infinitesimal rigidity result for the induced metrics (lemma 1.7) happens to be sufficient to obtain a satisfactory global result on the induced metrics.

Theorem 9.8. *Let S be a surface of genus $g \geq 2$, and let $N \geq 1$. For each complete, finite area hyperbolic metric h on S with N cusps, there is a unique ideal fuchsian hyperbolic manifold M such that the induced metric on each component of the boundary is h .*

The proof is in section 9.

1.5 Some hints on the techniques

The main idea behind the results obtained here is already a few years old: it is the use of the striking properties of the volume function on the space of hyperbolic structures. The first property is that the volume is a concave function when the space of hyperbolic structures is parametrized by the dihedral angles (this comes from the elementary lemma 2.5 concerning the ideal simplex) and the second is the Schläfli formula, which provides a fundamental link between the dihedral angles and the induced metric on ideal polyhedra, or on hyperbolic manifolds with ideal-like boundary.

Those ideas can be traced back to the treatment of the Andreiev theorem in [Thu97]; the existence of a "generating function" for the problem was then pointed out in [CdV91], and this function was identified as the volume in [Brä92] and [Riv94].

Following this path for hyperbolic manifolds with ideal-like boundary, however, leads to some technical difficulties. The first is that it is not clear how one can find an ideal triangulation of the manifold; by the way, a similar difficulty appears in other situations where one wants to deform hyperbolic structures, see e.g. [PP00]. This is treated in section 5, where it is shown that some finite cover of the manifolds

considered have an ideal triangulation. All the arguments can then be given in this finite cover, and an equivariance argument brings the result down to the manifolds we want to study.

Another technical point is to prove that the set of possible dihedral angle assignments (as it appears in theorem 8.15) is connected; this is necessary because the proofs rely on a deformation argument. The solution chosen here uses the fact that the conditions are "independent" on each of the connected components of the boundary. Moreover, if M has incompressible boundary, then the conditions on each connected component of ∂M is the same as for the corresponding fuchsian ideal polyhedra; so that the connectedness in the general case follows from the understanding of the ideal fuchsian polyhedra. So those fuchsian polyhedra are studied in section 4. The methods used there are quite different from those of the other sections; an existence and uniqueness result is proved for the third fundamental forms of fuchsian manifolds whose boundary locally looks like a compact polyhedron in H^3 , and from there one deduces an existence result for manifolds with ideal-like boundary by an approximation argument.

Although some other technically interesting details appear at different points in the paper, it does not seem necessary to describe them here — the reader will probably enjoy finding them as he reads along.

Even in the case of ideal polyhedra in H^3 , I think that the approach used here is different, and I believe slightly more direct, than the one used in previous papers — although no new result is achieved. Moreover, I hope that the methods used — as mentioned above, mostly developed earlier to study ideal polyhedra — could hint at some approaches to the questions concerning the convex cores of complete, convex co-compact hyperbolic manifolds. Of course many difficulties remain in this direction, although recent works of Bonahon [Bon98a, Bon98b] might provide useful tools.

Note that some of the results presented here — notably theorem 8.15 — have a non-empty intersection with some of the results obtained by Rivin in [Riv92]. The methods used, and their scope, however, are quite different.

1.6 What follows

Section 2 contains a reminder of the classical Schläfli formula, as well as some remarks and interpretations in terms of symplectic geometry. Section 3 then gives some basic definitions and results on the geometry of ideal hyperbolic manifolds.

The case of fuchsian manifolds is described in section 4, using methods that are much closer to those used in the classical theory of hyperbolic polyhedra (as developed in particular by Aleksandrov [Ale58]), and some methods from [LS00].

Section 5 deals with some technical questions on triangulations of ideal hyperbolic manifolds; the point is that, although I do not know how to construct a well-behaved triangulation of such a manifold, it is not too difficult to construct one on a finite cover, and this will be sufficient for this paper.

The geometrical constructions start in section 6, where more general hyperbolic structures on triangulated manifolds are investigated. The idea — which basically comes from earlier works, see [Thu97, CdV91, Brä92, Riv96] is to use variational properties of the volume functional on a larger class of hyperbolic structures to obtain hyperbolic metrics. This is continued in section 7, where first-order properties of the volume are described.

This leads in section 8 to results concerning dihedral angles, and in section 9 to results on induced metrics on the boundary. Some applications to circles packings are mentioned in section 10, and other considerations stand in the last section.

2 The Schläfli formula

We recall in this section the classical Schläfli formula. It is the main tool used in the sequel, so we also describe some interesting interpretations of it and some related properties of the volume of simplices.

We first state the Schläfli formula for compact polyhedra; for a proof, see e.g. [Mil94] or [Vin93].

Lemma 2.1. *Let (P_t) be a one-parameter family of convex polyhedra in H^3 . Let I be the set of its edges, $(L_i)_{i \in I}$ be its edge lengths, and $(\theta_i)_{i \in I}$ the corresponding (interior) dihedral angles. Then:*

$$dV = -\frac{1}{2} \sum_i L_i d\theta_i . \quad (1)$$

2.1 A symplectic viewpoint

There is an amusing symplectic interpretation of this formula. To explain it simply we choose an abstract polyhedron P_0 , with v vertices, e edges and f 2-faces, and consider only polyhedra with the same combinatorics. Call $\mathcal{L} := \mathbf{R}^e$ and $\Theta := (0, \pi)^e$ the sets containing the possible lengths and exterior dihedral angles, respectively, of hyperbolic polyhedra having the same combinatorial type as P_0 . Now consider the symplectic vector space $(\mathcal{L} \times \Theta, \omega)$, with $\omega := \sum_i dL_i \wedge d\theta_i$. Let $\mathcal{P}$ be the subset corresponding to the edge lengths and dihedral angles of convex polyhedra of the same combinatorial type as P_0 . It is well known, and not difficult to prove, that:

Proposition 2.2. *$\mathcal{P}$ is a submanifold (with boundary) of $\mathcal{L} \times \Theta$ of dimension e .*

We leave the proof to the reader, it uses the Euler formula and the fact that the number of constraints on the positions of the vertices is the weighted number of non-triangular faces.

The Schläfli formula is essentially equivalent to the following:

Corollary to the Schläfli formula. *$\mathcal{P}$ is lagrangian in $(\mathcal{L} \times \Theta, \omega)$.*

Proof. Define the 1-form:

$$\beta := \sum_i L_i d\theta_i.$$

Then $d\beta = \omega$. Since $\beta|_{\mathcal{P}} = dV|_{\mathcal{P}}$, $d\beta|_{\mathcal{P}} = 0$ so that ω vanishes on $\mathcal{P}$. □

2.2 Ideal polyhedra

Now let $\mathcal{P}_\infty$ be the set of ideal polyhedra having the combinatorial type of P_0 ; they are obtained by letting the vertices go to infinity – this is possible for many combinatorial types. A basic remark is that those polyhedra still have finite volume. Their edge lengths, however, are not defined as such (they are infinite). To define an analog of the edge lengths for ideal polyhedra, we choose for each vertex V a horosphere H_V “centered” on V , that is, a level set for the Busemann function associated to V . We then define the length of the edge joining 2 vertices V and W as the distance, along the edge, between H_V and H_W ; we use the signed length, so that the length is negative if the horospheres overlap. The set of edge lengths of the elements of $\mathcal{P}_\infty$ is thus defined up to the addition of a constant for each vertex, and is contained in the set $\mathcal{L}_\infty := \mathbf{R}^e / \mathbf{R}^v$.

On the other hand, the dihedral angles of the ideal polyhedra are constrained. That is because the link of each vertex is a Euclidean polygon, so the sum of its exterior angles is 2π , so that the sum of the exterior dihedral angles of the edges containing a given vertex of an ideal polyhedron is always 2π . Therefore, the set of dihedral angles of the polyhedra in $\mathcal{P}_\infty$ stays in a $(e - v)$ -dimensional space Θ_∞ .

The amusing fact that we announced is then:

Remark 2.3. *1. $\mathcal{L}_\infty \times \Theta_\infty$ is obtained from $\mathcal{L} \times \Theta$ by symplectic reduction, and it therefore still carries a symplectic form ω_∞ ;*

2. the Schläfli formula (1) still has a meaning, and still holds, for $\mathcal{P}_\infty$;

3. it still implies that $\mathcal{P}_\infty$ is a lagrangian submanifold of $(\mathcal{L}_\infty \times \Theta_\infty, \omega_\infty)$.

Proof. The second point is probably classical, and we leave it to the reader. For the first point, consider the group $G := \mathbf{R}^v$, and its action ϕ on $\mathcal{L} \times \Theta$ by:

$$\begin{aligned} \phi : \quad G = \mathbf{R}^v \times (\mathcal{L} \times \Theta) &\rightarrow \mathcal{L} \times \Theta \\ ((\alpha_j)_{j=1, \dots, v}, (L_i, \theta_i)_{i=1, \dots, e}) &\mapsto (L_i + \alpha_{i_+} + \alpha_{i_-}, \theta_i)_{i=1, \dots, e} \end{aligned}$$

where i_+ and i_- are the two ends of the edge i .

It is quite easy to check that ϕ has a moment map μ defined by:

$$\begin{aligned} \mu: \mathcal{L} \times \Theta &\rightarrow \mathcal{G}^* = \mathbf{R}^v \\ (L_i, \theta_i) &\mapsto \left(\sum_{k \in E_j} \theta_k \right)_{j=1, \dots, v} \end{aligned}$$

where E_j is the set of edges containing a vertex j .

Then $\mathcal{L}_\infty \times \Theta_\infty \simeq \mu^{-1}(2\pi, \dots, 2\pi)/G$, where G acts by ϕ , so that $\mathcal{L}_\infty \times \Theta_\infty$ is obtained by symplectic reduction from $\mathcal{L} \times \Theta$ as announced, with a symplectic form ω_∞ .

For the last point note that β determines a well-defined 1-form β_∞ on $\mathcal{L}_\infty \times \Theta_\infty$, so again $\omega_\infty = d\beta_\infty$. But the Schläfli formula for ideal polyhedra shows that β_∞ vanishes on $\mathcal{P}_\infty$, which is thus lagrangian. $\square$

2.3 The volume function

We will also need some elementary and well-known properties of the volume of hyperbolic simplices, which we recall here for the reader's convenience. More details can be found for instance in [Thu97], chapter 7.

Definition 2.4. *The Lobachevsky function is defined as:*

$$\Lambda(\theta) := - \int_0^\theta \log |2 \sin u| du .$$

Now recall that there is a 2-parameter family of ideal simplices in H^3 (up to global isometries), which can be parametrized for instance by the complex cross-product of the four vertices in $\partial_\infty H^3 \simeq \mathbf{CP}^1$. For each ideal simplex, the dihedral angles of two opposite edges are equal, and the sum of the exterior dihedral angles of the edges containing a given vertex is 2π . An ideal simplex is completely determined – again up to global isometry – by its three interior dihedral angles α, β and γ , under the condition that their sum is π .

The volume of an ideal simplex is given by a simple formula (see e.g. [Thu97], chapter 7):

Lemma 2.5. *The volume of the ideal simplex with dihedral angles α, β and γ is $\Lambda(\alpha) + \Lambda(\beta) + \Lambda(\gamma)$.*

As a consequence, one finds (see e.g. [Riv94]):

Corollary 2.6. *V is a concave function of the dihedral angles α, β, γ , which vanishes when one of the angles goes to 0.*

Proof. This is proved by an elementary computation of the Hessian of V . $\square$

Note that the Schläfli formula is not restricted to hyperbolic polyhedra; it is also valid in the other Riemannian or pseudo-Riemannian space-forms (see [SP00]) and also has an interesting extension to the setting of Einstein manifolds with boundary, see [RS00, RS99, Ber]. I do not know whether any analog of the methods described in this paper – where the volume is used as a "generating function" to prove geometric results – can be found in this more general context.

There is another, completely elementary proof of corollary 2.6, which uses the Schläfli formula instead of explicit computations of the volume in terms of the Lobachevsky function. It is also more general. The key point is that, according to the Schläfli formula, the infinitesimal deformations of an ideal simplex which do not change the lengths of the edges (i.e. the isometric deformations) are exactly the deformations which are in the kernel of the Hessian of the volume, seen as a function of the dihedral angles. Now it is easy to see — and classical — that ideal simplices are rigid, i.e. they have no non-trivial infinitesimal isometric deformation. Therefore, the Hessian of the volume has constant signature. But an elementary argument (with the Schläfli formula) shows that the regular simplex has maximal volume, so that V is a concave function.

This line of reasoning can of course be used in different contexts; for instance, it shows that the signature of the Hessian of the volume, as a function of the dihedral angles, has constant signature on the space of compact hyperbolic simplices, since they are also known to be rigid.

3 Convex hulls of ideal points

The goal of this paper is to understand hyperbolic manifolds whose boundaries look locally like ideal polyhedra. We will introduce a class of manifolds with boundary obtained by taking the convex hull of a finite number of ideal points in a complete, convex co-compact hyperbolic manifold. The main point is that the class of those manifolds with boundary separates into two sub-class, according to whether their "convex core" intersects their boundary or not. We will then indicate why both cases actually exist in a non-trivial way, since it is not completely obvious at first sight.

3.1 Some definitions

We first define a notion of convexity — it is classical but other definitions are sometimes used.

Definition 3.1. *Let M be a hyperbolic manifold. We say that a subset $C \subset M$ is **convex** if, for any points x and y in C , any geodesic segment in M with endpoints x and y remains in C . For any subset E of $M \cup \partial_\infty M$, the **convex hull** of E is the smallest convex set containing E .*

Definition 3.2. *M is a **hyperbolic manifold with horns** if there is a complete co-compact hyperbolic manifold N and a finite family $x_1, \dots, x_p$ of points in $\partial_\infty N$ such that M is isometric to the convex hull in N of $\{x_1, \dots, x_p\}$.*

For instance, the interior of an ideal polyhedron in H^3 is a hyperbolic manifold with horns. So is the convex core of a convex co-compact hyperbolic manifold; in this case $p = 0$.

Remark 3.3. *Let M be a compact hyperbolic manifold with convex boundary, such that the induced metric on the boundary is a complete hyperbolic metric of finite area on the boundary minus a finite number of points. Then M is a hyperbolic manifold with horns.*

Proof. Since M is compact with convex boundary it is isometric to a subset of a complete convex co-compact hyperbolic manifold N . Let $\tilde{M}$ be the universal cover of M , then $\partial\tilde{M}$ is a convex surface in H^3 and its induced metric is hyperbolic; it is therefore the convex hull in H^3 of its boundary points in $\partial_\infty H^3$. Taking the quotient by $\pi_1 M$, we see that ∂M is the convex hull of its boundary points in $\partial_\infty M$. Those points clearly correspond to the cusps of ∂M , so M is the convex hull in $H^3/\pi_1 M$ of a finite number of ideal points. $\square$

Consider a hyperbolic metric on M for which ∂M is convex. There is then a unique convex co-compact hyperbolic 3-manifold N in which M admits an isometric embedding which is surjective on the π_1 ; we call it the **extension** of M , and denote it by $E(M)$. The convex core of $E(M)$ can be defined as the smallest convex subset of $E(M)$, so it is contained in M ; we will denote it by $C(M)$.

Definition 3.4. *Let M be a hyperbolic manifold with horns. M is an **ideal hyperbolic manifold** if $C(M) \cap \partial M = \emptyset$, otherwise M is a **bent hyperbolic manifold**.*

For instance, the interiors of ideal polyhedra in H^3 are of the ideal kind, while the convex cores of convex co-compact manifolds are of the bent type. Most of what follows concerns ideal hyperbolic manifolds only, while the bent case appears as the problem lurking in the background. The relationship between the two will be explored after the next subsection, using the tools that it contains.

3.2 Relation with circle packings

To understand the behavior of the boundary of a hyperbolic manifold with horns M , it is relevant to consider its universal cover $\tilde{M}$, which can naturally be identified with a convex subset of H^3 . The vertices of $\tilde{M}$ correspond to ideal points in H^3 , i.e. to points in $\partial_\infty H^3$. The combinatorics of the faces of $\partial\tilde{M}$ is then described in terms of Delaunay cellulations of the sphere.

Definition 3.5. *Let σ be a cellulation of S^2 . σ is **Delaunay** if, for each cell c of σ :*

1. *the vertices of c are co-cyclic, i.e. lie on a circle $C(c)$ in S^2 ;*

2. one of the closed disks bounded by $C(c)$ contains no other vertex of σ .

Note that this definition involves only the vertices of σ , not its edges. It depends on the Möbius structure of S^2 , but not on its metric structure.

Lemma 3.6. *Let E be a discrete subset of S^2 . There is a unique maximal Delaunay cellulation of a subset of S^2 whose vertices are the elements of E . Its combinatorics is that of the convex hull of E in $\mathbf{R}^3$.*

The proof is elementary; the circles appearing in the definition of a Delaunay cellulation are the boundaries at infinity of support planes of the convex hull of E .

Let M be a hyperbolic manifold with horns. By definition, M is the convex hull of a finite set of ideal points $S = \{x_1, \dots, x_p\} \subset \partial_\infty E(M)$. Consider the universal cover $\tilde{E}(M) = H^3$ of $E(M)$; S lifts to a set of points $\tilde{S}$ which is invariant under the action of $\pi_1(M)$, so that the accumulation set of $\tilde{S}$ is the limit set $\Lambda \subset S^2$ of the action of $\pi_1 M$ on H^3 .

Consider a cellulation σ of a subset Ω of S^2 ; we consider it as of a partly geometric and partly combinatorial nature, with fixed vertices in S^2 but edges defined up to isotopy.

The following statement is a consequence of lemma 3.6.

Proposition 3.7. *There is a unique Delaunay cellulation of $S^2 \setminus \Lambda$ with vertices the elements of $\tilde{S}$; it is obtained as the combinatorial structure of the convex hulls of $\tilde{S}$ in H^3 .*

The circles appearing in the definition of a Delaunay cellulation are simply the traces on $\partial_\infty H^3$ of the 2-planes which are the faces of the convex hull of $\tilde{S}$. Moreover, the dihedral angles between those faces are the angles between the corresponding circles in $S^2 = \partial_\infty H^3$. Thus, dihedral angle questions on ideal hyperbolic manifolds can be translated as questions on configurations of circles in S^2 having given angles (and invariant under group actions). A precise relationship with circle packings can be obtained using an idea of Thurston; this is recalled in section 10.

We can now show that ideal hyperbolic manifolds have a simple description in terms of the properties of their boundary. Of course this description does not apply to bent hyperbolic manifolds.

Property 3.8. *Let M be an ideal hyperbolic manifold. Then its boundary ∂M is the union of a finite number of 2-faces, which are ideal polygons with a finite number of edges in totally geodesic 2-planes, and which intersect along geodesics.*

Proof. Since $E(M)$ is convex co-compact, its convex core $C(M)$ is compact. By definition, ∂M does not intersect $C(M)$, so $d(\partial M, C(M)) > 0$. Therefore, each face of ∂M remains at a positive distance from $C(M)$. Thus each face of ∂M remains in a compact subset of the complement of the limit set Λ of the action of $\pi_1 M$ on H^3 . Therefore, the circles in $S^2 = \partial_\infty H^3$ which are the boundary at infinity of those planes are not tangent to Λ .

Let Ω be a fundamental domain with a compact closure in $S^2 \setminus \Lambda$ for the action of $\pi_1 M$. Ω is contained in a compact subset $K \subset S^2 \setminus \Lambda$, so it intersects a finite number of those circles; therefore, ∂M has a finite number of faces. The proof of the property follows. $\square$

3.3 Existence of the bent case

Let N be a complete, convex co-compact hyperbolic 3-manifold. Let $x_1, \dots, x_p$ be a family of points in $\partial_\infty N$. If one of the connected component $\partial_0 N$ of $\partial_\infty N$ does not contain any of the x_i , it is not difficult to see that the smallest convex subset of N containing the x_i will be of the bent type; indeed, its boundary will contain the component of the boundary of the convex core of N facing $\partial_0 N$. We will sketch here an argument intended to convince the reader that there are other situations where bent manifolds appear.

A first remark is that a hyperbolic manifold with horns is bent if and only if, among the circles in $S^2 \setminus \Lambda$ which are the boundary at infinity of its faces, one has non-empty intersection with Λ . Indeed it was already proved in property 3.8 that this does not happen for an ideal hyperbolic manifold, while the converse is clear since, for such a circle, there should be a face of ∂M which is at distance 0 from $C(M)$.

Thus, to show the existence of a bent hyperbolic manifold, it is simplest to search for one among hyperbolic manifolds with horns with only one ideal point in each boundary component (those manifolds

could legitimately be called "unicorn manifolds"). Moreover we can restrict our attention to e.g. the quasi-fuchsian case — we will see below that fuchsian hyperbolic manifolds can not be bent.

Fix one of the boundary components of M , say $\partial_1 M$, and let $x_1 \in \partial_\infty M$ be the corresponding ideal point. One of the faces of $\partial_1 M$ has non-empty intersection with Λ if and only if there is a circle C_0 in S^2 , whose interior is in $S^2 \setminus \Lambda$, which intersects Λ , and whose interior contains no point of the orbit $(\pi_1 M).x_1$.

So, to prove the existence of a bent hyperbolic manifold, it is enough to prove the existence of a closed disk C_0 which:

1. has its interior in $S^2 \setminus \Lambda$.
2. intersects Λ .
3. contains no fundamental domain for the action of $\pi_1 M$ on S^2 .

If such a circle exists, there will be a point x_1 whose orbit $(\pi_1 M).x_1$ does not intersect the interior of C_0 .

To simplify a little the picture, suppose that M has a convex core whose pleating locus contains a closed geodesic γ , with a non-zero pleating angle. In H^3 , γ lifts to a geodesic $\tilde{\gamma}$ with endpoints $p_1, p_2 \in \Lambda$; both p_1 and p_2 then correspond to "spikes" of Λ . Moreover, $\tilde{\gamma}$ lies in the boundary of $C(\tilde{M})$, so there is a support plane P of $C(\tilde{M})$ along $\tilde{\gamma}$. Let C_0 be the corresponding disk in $S^2 = \partial_\infty H^3$. By construction, the interior of C_0 does not intersect Λ (while $C_0 \cap \Lambda \supset \{p_1, p_2\}$).

Call Ω_1 the connected component of $S^2 \setminus \Lambda$ which corresponds to $\partial_1 M$. Let g_1 be the hyperbolic metric on Ω_1 , conformal to the canonical metric g_0 on S^2 and invariant under the action of $\pi_1 M$. Then, by a classical result of conformal geometry (see e.g. [Ahl66]), the conformal factor between g_0 and g_1 is bounded between c/r and C/r , where c and C are two positive constants and r is the distance to Λ in the metric g_0 .

Therefore, a simple computation shows that the interior of C_0 is contained in a neighborhood of a geodesic in (Ω_1, g_1) , and also — by taking the quotient by $\pi_1 M$ — in $(\partial_1 M, g_1)$. Thus, after taking a finite cover of $\partial_1 M$, the interior of C_0 contains no fundamental domain for the action of $\pi_1 M$. Taking the corresponding finite cover $\tilde{M}$ of M , we see that $\tilde{M}$ is a bent hyperbolic manifold.

On the other hand, this argument directly shows that bending laminations can not occur in the case of fuchsian hyperbolic manifolds.

Remark 3.9. *Fuchsian hyperbolic manifolds with horns are ideal hyperbolic manifolds.*

Proof. When M is a fuchsian manifold with horns, Λ is a circle in $S^2 = \partial_\infty H^3$. Circles tangent to Λ have interiors which are isometric — for the hyperbolic metric on the interior of Λ — to horoballs in H^2 . Therefore they always contain fundamental domains for any co-compact action on H^2 . $\square$

3.4 Necessary conditions on convex surfaces

We recall here some well-known properties of the induced metric and third fundamental form of convex surfaces in H^3 ; they are classical except for the length condition on the geodesics of the third fundamental form, which was understood more recently.

To understand the third fundamental form of non-smooth convex surfaces, it is helpful to know how the duality between H^3 and S_1^3 works, so we will recall it rapidly here; see e.g. [Thu97, RH93, Sch98a] for more details. Both spaces can be seen as quadrics in the Minkowski 4-space $\mathbf{R}_1^4$, with the induced metric:

$$H^3 = \{x \in \mathbf{R}_1^4 \mid \langle x, x \rangle = -1 \wedge x_0 > 0\},$$

$$S_1^3 = \{x \in \mathbf{R}_1^4 \mid \langle x, x \rangle = 1\}.$$

For $x \in H^3$, let D be the line in $\mathbf{R}_1^4$ going through 0 and x , and let $D^\perp$ be its orthogonal for the Minkowski inner product, so that $D^\perp$ is a space-like plane. Then define the dual x^* of x as the intersection of $D^\perp$ with S_1^3 , which is a totally geodesic space-like plane in S_1^3 . Similarly, the dual of a point in S_1^3 is an oriented plane in H^3 . The dual of a convex polyhedron P in H^3 is the polyhedron in S_1^3 whose vertices are the duals of the faces of P , and whose faces are the duals of the vertices of P (note that there are other approaches of this duality, which might actually be more illuminating; see e.g. [Sch98a]).

Given a locally convex surface S in H^3 , we can define its dual S^* as the set of points in S_1^3 which are duals of an (oriented) support plane to S . It happens to be another locally convex surface, which is not necessarily smooth. If S is smooth, then S^* is smooth when S is locally strictly convex.

Property 3.10. *Let S be a smooth locally strictly convex surface in H^3 ; its third fundamental form is the induced metric on the dual surface.*

It is tempting to speak of the "third fundamental form" of a non-smooth surface in the sense "the metric induced on its dual". Some care is needed, however. For instance, if S is a connected component of the boundary of the convex core of a convex co-compact manifold, its dual is a graph; more generally, when S is a convex, developable surface, like the boundary of a horned hyperbolic manifold, its dual is one-dimensional.

On the other hand, this notion of third fundamental form works perfectly well for compact polyhedra in H^3 — and thus also for objects which locally look like them. For ideal polyhedra it also works quite well; in some cases (see below in part 3.5) it is helpful to "glue" a hemisphere in each of the length 2π circles corresponding to the ideal vertices.

Lemma 3.11. *Let S be a smooth (resp. polyhedral) locally convex surface in H^3 , and let I and III be its induced metric and third fundamental forms respectively. Then:*

1. *I has curvature $K \geq -1$ (resp. has curvature -1 , except at the vertices, where the singular curvature is positive);*
2. *III has curvature $K \leq 1$ (resp. has curvature 1 , except at the dual vertices, where the singular curvature is negative);*
3. *the closed geodesics of III have length $L \geq 2\pi$.*

In statements (1) and (2), the equality is attained only in the degenerate cases, i.e. when the surface is not locally strictly convex. In statement (3) it also corresponds to a very degenerate case, as we will see below.

Proof. The first and second point are consequences of the Gauss formula in the smooth case and can be checked locally in the polyhedral cases. For the last point the reader is referred to [RH93, CD95, BB02, Sch98a, Sch01a] for different approaches of the polyhedral case, and e.g. to [Sch96] for the smooth case. $\square$

We will also prove here lemma 1.6. The proof is based on a number of propositions. In all this subsection we consider a cellulation σ of ∂M , along with a function w on the edges of σ verifying the hypothesis of theorem 8.15.

Proposition 3.12. *No 2-cell of σ can have two edges sent to the same segment in $\partial\tilde{M}$. The intersection of two distinct 2-cells in $\partial\tilde{M}$ can not contain more than one 1-cell.*

Proposition 3.13. *No 2-cell can have two vertices sent to the same point in $\partial\tilde{M}$.*

Proposition 3.14. *The intersection of two 2-cells in $\partial\tilde{M}$ can not contain both a vertex and an edge (which are disjoint).*

Proposition 3.15. *The intersection of two 2-cells in $\partial\tilde{M}$ can not contain two vertices.*

The proof of lemma 1.6 clearly follows from those propositions.

Proof of proposition 3.12. Suppose that two edges of a 2-cell C are sent to the same segment in $\partial\tilde{M}$. This segment would then constitute a circuit in the 1-skeleton of σ , on which the sum of the values of w is strictly less than π ; this would contradict the hypothesis of theorem 8.15.

Similarly, if two 2-cells C and C' in $\partial\tilde{M}$ have two edges in common, then they constitute a circuit on which the sum of the values of w is strictly less than 2π . $\square$

Proof of proposition 3.13. Let C be a 2-cell having two vertices which are sent to the same point x_0 in $\partial\tilde{M}$. Since the edges of C are sent to distinct segments by the proposition 3.12, C separates $\partial\tilde{M}$ in two parts, one compact and the other non-compact, whose closure intersect at x_0 .

Let c be the elementary circuit made of the edges adjacent to x_0 . Then c is the union of two sequences of edges, one, say c_i , made of the edges contained in the closure of the compact domain of the complement of C , and another, say c_o , made of the other edges. Both sequences constitute a non-elementary circuit, and the sum of the values of w is less than 2π on both. This again contradicts the hypothesis of theorem 8.15. $\square$

Proof of proposition 3.14. Suppose that two cells C and C' have in common both a vertex, x_0 , and an edge e . Let c be the elementary circuit made of the edges adjacent to x_0 . Again, c can be written as the union of c_i and c_o , where c_i is the sequence of edges contained in the bounded domain in the complement of $C \cup C'$. Adding e to either c or c' leads to a non-elementary circuit. But, according to the hypothesis of theorem 8.15 the sum of the values of w on the edges of c is 2π , so that the sum of the values of w has to be less than 2π either on the edges of c_i and e , or on the edges of c_o and e . $\square$

Proof of proposition 3.15. Suppose now that C and C' share two vertices x_0 and x_1 . Let c be the elementary circuit made of the edges adjacent to x_0 , and let c' be the elementary circuit made of the edges adjacent to x_1 . Again, both c and c' decompose into two sequences of edges, those in the closure of the bounded domain in the complement of $C \cup C'$ (call them c_i and c'_i , respectively) and the others (let them be c_o and c'_o , respectively).

Then $c_i \cup c'_i$, $c_i \cup c'_o$, $c_o \cup c'_i$ and $c_o \cup c'_o$ each is a non-elementary circuit, while the sum of the values of w on at least one of them is clearly at most 2π , again contradicting the hypothesis of theorem 8.15. $\square$

3.5 Third fundamental form versus dihedral angles

One of the points which should be clear from the introduction is that we want to consider the manifolds with polyhedral boundary together with those having smooth boundary. It is in that respect necessary to understand what the relationship between the third fundamental form and the dihedral angles is. This should clear up how question concerning the third fundamental form — as in theorem 1.2 — are related to questions on the dihedral angles — as in theorem 8.15. We will work out the relationship here; it is mostly well-known, the most interesting case is that of ideal polyhedra.

First note that for manifolds with a boundary that looks locally like a compact hyperbolic polyhedron one can consider the universal cover of M , and then the surface $\tilde{S}^*$ in S_1^3 which is dual to the universal cover of a connected component of ∂M . $\tilde{S}^*$ is a polyhedral surface — locally like a convex space-like polyhedron in S_1^3 , so that it carries a spherical cone-metric with negative singular curvature at the singularities. The dihedral angles are then just the lengths of the dual edges.

This metric — which we will still call the third fundamental form of the boundary — lifts to a CAT(1) metric on the boundary of the universal cover of M . Indeed, this splits into a local curvature condition — which is satisfied by the local convexity, because the singular curvature is negative at each vertex — and a global condition on the length of the closed geodesics, which is also true here because of lemma 3.11.

A rather important point is that, while knowing the dual metric (and the combinatorics of the polyhedral surface) determines the dihedral angles, the converse is not true — the dihedral angles determine the length of the edges of the dual surface, but it does not determine the shape of the dual faces with more than 4 edges. This is already the source of interesting questions for convex polyhedra in H^3 ; for instance, it is still an open problem to know whether a convex hyperbolic polyhedron can be infinitesimally deformed without changing its dihedral angles, see e.g. [Sch00], or [Sto68] for an analogous (and also open) problem in the Euclidean case.

For ideal polyhedra, there are two related ways of defining the third fundamental form. If one considers the dual surface to the universal cover of one of the components of the boundary, one obtains the induced metric on the dual of a pleated surface, which is a tree. The third fundamental form therefore reduces to the lengths of the edges of a graph. The length of the edges are the (exterior) dihedral angles of the corresponding edges of the ideal polyhedron.

To each ideal vertex corresponds a face of the dual graph, with the sum of the edge lengths equal to 2π (since the sum of the exterior dihedral angles at an ideal vertex is 2π). One can glue in each of those faces a hemisphere (with its canonical metric). The result is a metric space on ∂M , which obviously has negative singular curvature at its singular points, because the singular points correspond to the vertices of the graph, and the total angle around those points is π times the number of faces. We call $\mathbb{III}$ the corresponding metric on ∂M . Note that $\mathbb{III}$ is the “natural” third fundamental form of ∂M for instance in a limit sense, as follows:

Property 3.16. *Let $(\Omega_n)_{n \in \mathbb{N}}$ be an increasing sequence of open subsets of M with smooth, convex boundary, such that $\cup_n \Omega_n = M$. Then the third fundamental forms of $\partial \Omega_n$ converge to $\mathbb{III}$.*

We leave the proof to the reader.

The third fundamental form defined in this way has the important properties below. The second strongly contrasts with the situation for compact polyhedra.

Property 3.17. *1. $\mathbb{III}$ lifts to a CAT(1) metric on each boundary component of the universal cover of M .*

2. There is a simple way to recover the cellulation of ∂M from $\mathbb{III}$, and therefore also the dihedral angles.

Proof. The first point is again a consequence of lemma 3.11.

For the second point, note that the dihedral angles of ideal polyhedra are in $(0, \pi)$, and so are the edge lengths of the dual surfaces. Now any geodesic segment that enters a hemisphere can exit it only after a path of length π . Since the dihedral angles are less than π , the edges of the dual cellulation can not enter the hemispheres. On the other hand all the segments in the boundary of the hemispheres must be edges, and this recovers the dual cellulation, and thus also the cellulation of ∂M . $\square$

4 The fuchsian case

We will investigate in this section some properties of manifolds with polyhedral boundary in the fuchsian case; that is, we consider a metric g on M such that ∂M is polyhedral, with an isometric involution s of (M, g) which fixes a compact surface. We also suppose that ∂M have two connected components, which are exchanged by s . We refer the reader to [Lei01a, Lei01b] for some recent results on those manifolds.

Another way to consider such manifolds is to take the universal cover S of one the two connected components of ∂M ; it is a convex surface in H^3 , which moreover is equivariant under the action of a surface group fixing a totally geodesic 2-plane.

We will first prove an existence and uniqueness result for the third fundamental forms of such surfaces, in the case where they locally look like a compact (rather than ideal) polyhedron in H^3 . This is done using a deformation argument as in the case of hyperbolic polyhedra (see [Ale58]), and is also similar to what can be done in the smooth case [LS00].

This first result will then be used to prove a similar existence result for the dihedral angles of surfaces which locally look like an ideal polyhedron; the main idea is to approximate this case by the previous one, using a compactness result to prevent degeneracies from occurring. The proof will also give a result concerning the manifolds with a polyhedral boundary with some ideal and some “non-ideal” vertices.

The result which is obtained in this way — or at least its ideal part — might look like a weak and partial version of some of the results stated in the introduction. It is proved in a very different way, however, and turns out to be necessary for the more general cases, because it implies a technical statement — on the connectedness of some spaces of metrics on the boundary — which I do not know how to obtain directly.

Mathias Rousset [Rou02] has recently achieved another related result, concerning the dihedral angles of fuchsian hyperideal polyhedra, which includes the case of ideal fuchsian polyhedra. He reduces the study of hyperideal polyhedra to that of finite polyhedra.

In all this section, we fix a surface S of genus $g \geq 2$; we will be interested in equivariant embeddings of S , or, in other terms, in fuchsian hyperbolic metrics on $S \times [-1, 1]$.

4.1 Infinitesimal rigidity of finite fuchsian polyhedra

We first consider equivariant polyhedra which look locally like compact polyhedra. First recall the definition of a polyhedral embedding.

Definition 4.1. A **polyhedral embedding** of a surface S into H^3 (resp. $\mathbf{R}_1^3$) is a couple (σ, ϕ) , where σ is a cellulation of S and ϕ is a map from S to H^3 (resp. $\mathbf{R}_1^3$) which:

- is injective.
- sends each edge of σ to a segment in H^3 (resp. $\mathbf{R}_1^3$).
- sends each 2-face of σ to the interior of a compact, convex polygon in a totally geodesic 2-plane in H^3 (resp. $\mathbf{R}_1^3$).
- is locally convex at each vertex.

The last condition is not always necessary; it is included here since all the polyhedral objects that we will consider are convex.

Definition 4.2. A **finite equivariant polyhedron** is a couple (ϕ, ρ) , where ϕ is a polyhedral embedding of the universal cover of a surface S into H^3 and ρ is a group morphism from $\pi_1 S$ into $\text{Isom}(H^3)$, such that:

$$\forall x \in \tilde{S}, \forall \gamma \in \pi_1 S, \phi(\gamma x) = \rho(\gamma)\phi(x) .$$

We are specially interested in the equivariant polyhedra which are the boundary of the universal covers of the fuchsian hyperbolic manifolds mentioned above.

Definition 4.3. A finite equivariant polyhedron (ϕ, ρ) is **fuchsian** if $\rho(\pi_1 S)$ is contained in the subgroup of elements which leave invariant a given plane $P \subset H^3$.

The definition of ideal equivariant polyhedra, and of semi-ideal equivariant polyhedra, is similar. We first define the notion of polyhedral map.

Definition 4.4. An **ideal polyhedral embedding** of a surface S into H^3 is a couple (σ, ϕ) , where σ is a cellulation of S and ϕ is a map from S to $H^3 \cup \partial_\infty H^3$ which:

- is injective.
- sends each vertex of σ to an ideal point (in $\partial_\infty H^3$).
- sends each edge of σ to a geodesic in H^3 , which connects the ideal points corresponding to the vertices.
- sends each 2-face of σ to the interior of an ideal convex polygon in a totally geodesic 2-plane in H^3 .
- is locally convex .

A **semi-ideal polyhedral embedding** is defined likewise, except that the vertices can be sent either to ideal points or to “usual” points of H^3 .

Definition 4.5. An **ideal equivariant polyhedron** is a couple (ϕ, ρ) where ϕ is an ideal polyhedral embedding of the universal cover of a surface S into H^3 and ρ is a group morphism from $\pi_1 S$ into $\text{Isom}(H^3)$, such that:

$$\forall x \in \tilde{S}, \forall \gamma \in \pi_1 S, \phi(\gamma x) = \rho(\gamma)\phi(x) .$$

A **semi-ideal equivariant polyhedron** is defined in the same way, but with ϕ a semi-ideal polyhedral embedding.

Note that the induced metric on equivariant (semi-)ideal polyhedra is defined not only on the universal cover of the underlying surface, but also, by equivariance, on the quotient surface. For ideal equivariant polyhedra it is a hyperbolic metric with cusps corresponding to the vertices (i.e. a hyperbolic metric of finite area on the quotient surface S). For semi-ideal polyhedra, the induced metric is a cone-metric, with singular points corresponding to the vertices which are not ideal. It also has cusps corresponding to the ideal vertices.

Definition 4.6. An ideal (resp. semi-ideal) equivariant polyhedron (ϕ, ρ) is **fuchsian** if $\rho(\pi_1 S)$ is contained in the subgroup of elements which leaves invariant a given plane $P \subset H^3$.

The first result we need to mention is an infinitesimal rigidity result for finite equivariant polyhedral embeddings in Minkowski 3-space.

Lemma 4.7. Let (ϕ, ρ) be a convex equivariant space-like polyhedron in $\mathbf{R}_1^3$, such that the representation ρ fixes the origin. There is no non-trivial infinitesimal deformation of (ϕ, ρ) , preserving the condition that ρ fixes the origin, which does not change the induced metric by ϕ to the first order.

This statement can be found as theorem B in Igor Iskhakov's thesis [Isk00], where it is proved by an extension of Cauchy's ideas on the rigidity of polyhedra (see [Cau13, Sto68]) to surfaces of genus $g \geq 2$.

I guess that an alternate proof could be given, along the approach given in [LS00] for smooth surfaces; the key point would be to replace the integration by part which works for the smooth case by a discrete version.

There is yet another way to prove this infinitesimal rigidity result, as well as the infinitesimal rigidity in [LS00], and more general results concerning for instance hyperideal fuchsian polyhedra. It is based on the Pogorelov map, which is also used for instance in [Sch02a], to bring the problem in $\mathbf{R}^3$. There, the infinitesimal rigidity can be proved using the elementary fact that, if an infinitesimal deformation of a (smooth or polyhedral) convex surface is isometric, then the graph of its coordinates are saddle surfaces.

As a consequence, we find that the same result holds in the de Sitter space:

Lemma 4.8. Let (ϕ, ρ) be a convex equivariant space-like polyhedron in S_1^3 , such that the representation ρ fixes a point x_0 . There is no non-trivial infinitesimal deformation of (ϕ, ρ) , preserving the condition that ρ fixes x_0 , which does not change the induced metric by ϕ to the first order.

The proof of lemma 4.8 from lemma 4.7 is related to a remarkable trick invented by Pogorelov [Pog73], which allows one to take an infinitesimal rigidity problem from a space-form to another. The crucial point is that this can be done in this setting, i.e. for equivariant objects when the representation fixes a point, as shown in [LS00] for smooth surfaces. Moreover, the polyhedral case works just like the smooth case, details on this stand in [Sch00, Sch98a, Sch01a].

Note that the same could be done also in the anti-de Sitter space H_1^3 , and this is indeed done in [LS00] for smooth surfaces. We leave this point to the reader, however, since it will not be necessary below.

Using the duality between H^3 and the de Sitter space S_1^3 , we immediately find a translation of this lemma in terms of fuchsian surfaces in H^3 :

Corollary 4.9. Let (ϕ, ρ) be a convex equivariant polyhedron in H^3 , such that the representation ρ fixes a plane π_0 . There is no non-trivial infinitesimal deformation of (ϕ, ρ) , preserving the condition that ρ fixes π_0 , which does not change the third fundamental form of the image of ϕ to the first order.

4.2 Compactness of fuchsian polyhedra

The main technical tool of this subsection is a compactness result, which is necessary to obtain the existence and uniqueness result for fuchsian polyhedra explained above, and stated below as theorem 4.22. It is stated in a more general context, however, so as to be used also later in this section, to prove a result for ideal or semi-ideal fuchsian polyhedra.

We now fix two integers $N \geq 1$ and $g \geq 2$; g will be the genus of the surface S considered, and N will be the number of vertices of the polyhedral surfaces and the number of singular points of the metrics considered.

Definition 4.10. We call:

- $\mathcal{P}^C$ the set of finite fuchsian polyhedra of genus g with N vertices in H^3 .
- $\mathcal{P}^I$ the set of semi-ideal fuchsian polyhedra of genus g with N vertices in H^3 .

Definition 4.11. We call:

- $\mathcal{M}^C$ the set of spherical cone-metrics on S with N cone-points where the singular curvature is negative, and such that all contractible closed geodesics have length $L > 2\pi$ (up to isotopy).
- $\mathcal{M}^I$ the set of spherical cone-metrics with N singular points, where the singular curvature is negative, and such that contractible closed geodesics have length $L > 2\pi$, except when they bound a hemisphere (again up to isotopy).

It is clear (using lemma 3.11) that the third fundamental forms of the elements of $\mathcal{P}^C$ are in $\mathcal{M}^C$. Similarly, the third fundamental forms of the elements of $\mathcal{P}^I$ are in $\mathcal{M}^I$, after one makes a simple surgery: gluing a hemisphere on each of the circles, of length 2π , which correspond to the ideal vertices.

Lemma 4.12. *Let (ϕ_n, ρ_n) be a sequence of finite fuchsian polyhedral embeddings in S_1^3 , with representations ρ_n fixing a point x_0 . Let $(h_n) \in (\mathcal{M}^C)^{\mathbf{N}}$ be the induced metrics. Suppose that (h_n) converges, as $n \rightarrow \infty$, to a metric $h \in \mathcal{M}^I$. Then, after taking a subsequence and renormalizing, (ϕ_n, ρ_n) converges to a convex, fuchsian polyhedron (ϕ, ρ) . Moreover ρ fixes x_0 .*

In this statement, the “renormalization” is by composition on the left by an isometry.

Note that (ϕ, ρ) might have some faces which are tangent to the boundary at infinity of H^3 in the projective model of H^3 and S_1^3 . More precisely, this happens exactly when h has closed geodesics of length 2π bounding hemispheres, and the faces tangent to the boundary at infinity are precisely those hemispheres.

The proof of lemma 4.12 depends on the following propositions. The first is well known and its proof is easy, so we leave it to the reader.

Proposition 4.13. *Let H_0 be a space-like totally geodesic plane in S_1^3 . The function v on S_1^3 defined as the hyperbolic sine of the oriented distance to H_0 satisfies:*

$$\text{Hess}(v) = -vg_0 ,$$

where g_0 is the metric of S_1^3 .

This function will be used here — as in other similar problems, see e.g. [Sch96, LS00] — to control the lengths of closed geodesics of surfaces or the total extrinsic curvature of the surface along those geodesics.

The next proposition describes another way to write the canonical metric of the de Sitter space. The proof is again elementary.

Proposition 4.14. *The restriction to the future cone $C_+(x_0)$ of x_0 of the canonical metric of S_1^3 can be written as:*

$$v^2 g_{H^2} - \frac{dv^2}{1+v^2} ,$$

where g_{H^2} is the canonical metric on H^2 .

Proof. For $t > 0$, let Σ_t be the set of points x in the future cone of x_0 such that there is a time-like segment going from x_0 to x of length t . Then a simple computation shows that Σ_t has second fundamental form $\mathbb{I}_t = \coth(t)I_t$, where I_t is the induced metric on Σ_t . Moreover the surfaces Σ_t are equidistant, so that an integration shows that the induced metrics are:

$$I_t = \sinh^2(t)g_{H^2} ,$$

so that the metric on the future cone of x_0 can be written as:

$$\sinh^2(t)g_{H^2} - dt^2 .$$

The proposition follows by setting $v = \sinh(t)$. □

We then need a simple proposition about the solution of an ordinary differential inequality; it will be applied below to the function v restricted to geodesic segments in the metrics h_n .

Proposition 4.15. *For any $L > 0$ and $u_0 > 0$, there exists $c > 0$ such that if $u : [0, L] \rightarrow \mathbf{R}_+$ is a function which:*

- is smooth and satisfies $u'' = -u$ except at N singular points $x_1, \dots, x_N \in (0, L)$;
- has a positive jump in its derivative at the x_i ;
- is bounded from below by some constant $u_0 > 0$,

then:

$$\int_0^L \sqrt{\frac{1}{u^2} + \frac{u'^2}{u^2(1+u^2)}} ds \leq c .$$

Proof. First note that:

$$\begin{aligned} \sqrt{\frac{1}{u^2} + \frac{u'^2}{u^2(1+u^2)}} &\leq \frac{1}{|u|} + \frac{|u'|}{|u|\sqrt{1+u^2}} \\ &\leq \frac{1}{u_0} + \left| \left(\frac{1}{u} \right)' \right| . \end{aligned}$$

It is therefore enough to prove that the total variation over $[0, L]$ of $1/u$ is bounded from above by a constant.

Now let $0 \leq x_1 \leq \dots \leq x_p \leq L$ be the sequence of local minima of u , and let $0 \leq y_1 \leq \dots \leq y_q \leq L$ be its local maxima. The properties of u clearly imply that there is indeed a finite number of minima and maxima. Then, if y_j is a local maximum which immediately follows the local minimum x_i :

$$\int_{x_i}^{y_j} \left| \left(\frac{1}{u} \right)' \right| ds = \frac{1}{u(y_j)} - \frac{1}{u(x_i)} .$$

But, since y_j is a local maximum, and since u' has a positive jump at the singular points, $u'(y_j) = 0$, so that, for all $s \in [x_i, y_j]$:

$$u(y_j) \geq u(s) \geq u(y_j) \cos(y_j - s) .$$

We now consider two cases:

1. $|y_j - x_i| \geq \pi/4$. Then:

$$\left| \frac{1}{u(y_j)} - \frac{1}{u(x_i)} \right| \leq \frac{2}{u_0} .$$

2. $|y_j - x_i| \leq \pi/4$. Then:

$$\left| \frac{1}{u(y_j)} - \frac{1}{u(x_i)} \right| \leq \frac{1}{u(y_j)} \left(\frac{1}{\cos(y_j - x_i)} - 1 \right) \leq \frac{1}{u(y_j)} 4(y_j - x_i)^2 .$$

An elementary symmetry argument shows that the same estimates apply when x_j is a local minimum which immediately follows a local maximum y_i . As a consequence, we find that:

$$\int_0^L \left| \left(\frac{1}{u} \right)' \right| ds \leq \frac{2}{u_0} \frac{4L}{\pi} + \frac{4L^2}{u_0} ,$$

and the proposition follows. $\square$

We will also need the following more geometric estimate.

Proposition 4.16. *For any $r > 0$ and any integer $N > 0$, there exists $c > 0$ as follows. Let σ be a cellulation of the disk D with at most N vertices, and let ϕ be a polyhedral space-like embedding of D in S_1^3 with combinatorics given by σ , such that:*

- the boundary of $\phi(S)$ is convex for the metric induced by ϕ .

- the boundary is at distance at least r from the center x_0 in the induced metric.
- ϕ remains in the future cone $C_+(x_1)$ of a point x_1 .

Then the absolute value of the distance between $\phi(x_0)$ and x_1 is at least c .

Proof. Since $\phi(D)$ is space-like with convex boundary, it remains outside the future cone of each of its points, in particular outside the future cone of x_0 . Since it also remains in $C_+(x_1)$, it is not difficult to check that, if x_1 was too close to x_0 , $\phi(D)$ would have to remain in an arbitrarily small neighborhood of a cone; this is not possible for a polyhedral surface having a fixed number of singular points. $\square$

We will also need below a basic result in the theory of hyperbolic surfaces; see e.g. [FLP91] for a proof. Its content is that, to prevent a sequence of hyperbolic metrics from degenerating, one only needs to bound from above the lengths of a finite set of closed geodesics.

Lemma 4.17. *Let Σ be a surface of genus $g \geq 2$. There exists a finite subset E of $\pi_1\Sigma$ such that, for any $\epsilon > 0$, the set of hyperbolic metrics on Σ such that the closed geodesics corresponding to the elements of E have length at most $1/\epsilon$ is compact.*

We can now state a proposition showing that, with the hypothesis of lemma 4.12, the representations of the equivariant polyhedra (ϕ_n, ρ_n) do not diverge; the last part of the proof will be to show that, under the "length 2π " condition, isolated vertices can not escape to infinity.

Proposition 4.18. *In the setting of lemma 4.12, the sequence of representations (ρ_n) converges (after one takes a subsequence).*

Proof. First we fix an integer $n \in \mathbf{N}$, and consider an equivariant polyhedral embedding $\phi_n : \tilde{S} \rightarrow S_1^3$. Let x_n be a point of $\tilde{S}$ where the minimum of v is attained. Let $v_n := v \circ \phi_n$. Proposition 4.16 shows that $v_n(x_n)$ is bounded from below by a strictly positive constant. Let $E \subset \Gamma = \pi_1 S$ be a finite generating set, on which more details will be given below.

Let $\gamma \in E$. There is a minimal geodesic segment $c_{n,\gamma}$ going from x_n to γx_n in (S, h_n) . Since E is finite, and since $h_n \rightarrow h$ as $n \rightarrow \infty$, the length $L(c_{n,\gamma})$ of $c_{n,\gamma}$ is bounded from above by a constant L_0 for each $x \in E$ and each $n \in \mathbf{N}$.

Since $\phi_n(\tilde{S})$ is space-like and equivariant under the action of Γ , which fixes x_0 , $\phi_n(\tilde{S})$ remains in the future cone $C_+(x_0)$ of x_0 . Calling u_n the restriction of v to $c_{n,\gamma}$, proposition 4.13 shows that u_n satisfies $u_n'' = -u_n$, except when $c_{n,\gamma}$ crosses an edge of $\phi_n(\tilde{S})$, and then u_n' has a positive jump.

Proposition 4.14 implies that $C_+(x_0)$ has a natural submersion $\rho : C_+(x_0) \rightarrow H^2$, such that the restriction of ρ to each surface $\{v = \text{const}\}$ in $C_+(x_0)$ is a homothety. Let s be the length element induced by ϕ_n on S , and t be the length element of the hyperbolic metric induced on S by $\rho \circ \phi_n$. Proposition 4.14 then indicates that:

$$ds^2 = u_n^2 dt^2 - \frac{du_n^2}{1 + u_n^2} ,$$

$$dt^2 = \frac{ds^2}{u_n^2} + \frac{du_n^2}{u_n^2(1 + u_n^2)} ,$$

so that the length of the image of $c_{n,\gamma}$ by ρ is:

$$L_s(c_{n,\gamma}) = \int_0^{L(c_{n,\gamma})} \sqrt{\frac{1}{u_n^2} + \frac{u_n'^2}{u_n^2(1 + u_n^2)}} ds .$$

According to propositions 4.16, u_n is bounded from below by a constant $u_0 > 0$; proposition 4.15 therefore shows that, as $n \rightarrow \infty$, the lengths of the $s(c_{n,\gamma})$ remain bounded from above by a constant.

Now for each $n \in \mathbf{N}$, Γ acts on S_1^3 fixing x_0 , and therefore Γ has an action on $C_+(x_0)$ which leaves globally invariant all the surfaces $\{v = \text{const}\}$. Thus it acts by isometries on H^2 through ρ . The previous argument shows that the translation distance of each element of E remains bounded from above as $n \rightarrow \infty$. Lemma 4.17 then implies that the sequence ρ_n remains in a compact subset of Teichmüller space, and therefore that one of its subsequences converges. $\square$

To prove that the sequence of equivariant polyhedra (ϕ_n, ρ_n) actually converges — and not only the representations — it is helpful to consider a projective model of the part of the de Sitter space S_1^3 which stands on one side of a totally geodesic space-like plane. One can be constructed as follows. Remember that S_1^3 is isometric to a quadric in Minkowski 4-space with the induced metric:

$$S_1^3 \simeq \{x \in \mathbf{R}_1^4 | \langle x, x \rangle = 1\} .$$

Let P_0 be the affine hyperplane of equation $x_0 = 1$ in $\mathbf{R}_1^4$, and let:

$$S_{1,+}^3 := \{x \in S_1^3 | x_0 > 0\} .$$

There is a natural map from $S_{1,+}^3$ to P_0 sending a point $x \in S_{1,+}^3$ to the intersection with P_0 of the line going through 0 and x . By construction it is projective, i.e. it sends geodesics to geodesics; indeed, an elementary argument using the action of $\text{SO}(3,1)$ shows that the geodesics of $S_{1,+}^3$ are the intersections with $S_{1,+}^3$ of the 2-planes of $\mathbf{R}_1^4$ containing 0. Note that, since the $\phi_n(\tilde{S})$ remain in the future cone of a point, the projective model can be chosen so that a compact subset of $\mathbf{R}^3$ contains the image of the surfaces $\phi_n(\tilde{S})$ for each n .

Proposition 4.19. *After taking a subsequence and renormalizing, the sequence $(\phi_n(\tilde{S}))$ converges in the projective model described above to an equivariant polyhedron.*

Proof. This is a direct consequence of the convergence (after taking a subsequence) of the representations, as stated in proposition 4.18. Indeed, we can renormalize the sequence so that, for a given vertex $x \in \tilde{S}$, $\phi_n(x)$ is constant. Then an elementary compactness argument shows that, after taking a subsequence, the vertices adjacent to x also have converging images. Going to the vertices adjacent to those and applying the same compactness argument shows that they also have converging images, and this can be done until all vertices in a fundamental domain of $\tilde{S}$ have converging image. Proposition 4.18 then implies the result. $\square$

We now have to exclude some cases, corresponding to a limit metric h which has a closed geodesic of length 2π . A similar assertion was used — and stated in a more general setting, including higher dimensions — in [Sch98a, Sch01a].

Proposition 4.20. *Suppose there is a finite set of vertices which converge to the same point x_∞ in $\partial_\infty H^3$, while the other vertices do not. Then the limit metric h has a closed geodesic of length 2π which does not bound a hemisphere containing no vertex.*

Proof. We use proposition 4.19 and suppose that, in the projective model described above, the sequence $(\phi_n(\tilde{S}))$ converges to a limit P . Then $x_\infty \in P$, and, by convexity and the fact that the polyhedra $\phi_n(\tilde{S})$ remain in the exterior of the ball B^3 corresponding to H^3 , P also contains a neighborhood of x_∞ in the plane π_0 tangent to B^3 at x_∞ . More precisely, since P is a polyhedron, one of its faces is the interior of a convex polygon Q in π_0 .

We will show that the boundary polygon, ∂Q , has length 2π and is a geodesic for the limit induced metric h . The fact that it has length 2π is an elementary fact of Lorentz geometry, since it lies in a degenerate plane in S_1^3 . To show that it is a geodesic of h , we have to show that both sides of P are concave for h .

Consider first the interior of the polygon Q . Since it carries a degenerate metric, it is not difficult to realize that the metrics induced on the corresponding faces of the $\phi_n(\tilde{S})$ converge to the metric of a hemisphere. Therefore, this part of P is concave for h .

For the other side, simply note that, for each n , the metric h_n has a point of negative singular curvature at the vertices of ∂Q , i.e. the limit total angle at those vertices is at least 2π . Since the limit metric is isometric to a hemisphere in the interior of Q , it means that the limit total angle at each vertex of ∂Q of the complement of Q in P is at least π , i.e. that the complement of Q in P is concave for h , as needed. $\square$

Proof of lemma 4.12. Proposition 4.18 shows that the sequence of representations (ρ_n) converges (after taking a subsequence), while proposition 4.20 indicates that no vertex can escape to infinity. Therefore (ϕ_n, ρ_n) converges. $\square$

4.3 Induced metrics on finite polyhedra

We can now consider the maps $\Phi^C : \mathcal{P}^C \rightarrow \mathcal{M}^C$ and $\Phi^I : \mathcal{P}^I \rightarrow \mathcal{M}^I$ sending a fuchsian polyhedron to its third fundamental form. We will prove that Φ^C is a homeomorphism, and that Φ^I is bijective. To show that for Φ^C , we will apply a deformation method, which I believe was invented by Aleksandrov [Ale58] to study the induced metrics on hyperbolic polyhedra, although it was later used in many very different fields.

First, it is clear that choosing an element of $\mathcal{P}^C$ is equivalent to choosing a hyperbolic metric on S (or equivalently a fuchsian action of $\Gamma := \pi_1(S)$ on H^3) along with N points $x_1, \dots, x_N$ in H^3/Γ , under some conditions, i.e. that the x_i all lie on the boundary of their convex hull. Therefore, $\mathcal{P}^C$ is a connected manifold with boundary, of dimension $6g - 6 + 3N$.

It is also clear, using the results of Troyanov [Tro91], that choosing an element of $\mathcal{M}^C$ is the same as choosing an element of the Teichmüller space of S with N marked points, along with the singular curvature at each of the marked points. Thus $\mathcal{M}^C$ is also a manifold with boundary of dimension $6g - 6 + 3N$.

Moreover, corollary 4.9 shows that Φ^C is locally injective — and therefore a local diffeomorphism — between $\mathcal{P}^C$ and $\mathcal{M}^C$. To prove that it is a homeomorphism, we need to prove that Φ^C is proper (this is a consequence of lemma 4.12), that $\mathcal{P}^C$ is connected, and that $\mathcal{M}^C$ is simply connected.

The actual proof below is slightly more complicated than the outline here; since I do not know how to prove directly the connectedness of $\mathcal{M}^C$, we will use a trick invented in [Riv86, RH93]: one uses the connectedness of a space of smooth metrics — which is easy to prove — to check that two metrics g_0 and g_1 with N singular points can be connected by a path of metrics with at most N' singularities, where N' is a (large) integer depending on g_0 and g_1 .

Lemma 4.21. *1. Let g_0 and g_1 be elements of the space $\mathcal{M}(S, N)$ of spherical cone-metrics on S , with at most N singular points where the singular curvature is negative, and contractible closed geodesics of length $L > 2\pi$. There exists an integer N' depending on g_0 and g_1 such that g_0 and g_1 can be connected in $\mathcal{M}(S, N')$.*

2. Let $c : S^1 \rightarrow \mathcal{M}(S, N)$. There exists $N' \geq N$ and a disk $D \subset \mathcal{M}(S, N')$ with $\partial D = c(S^1)$.

Brief sketch of the proof . We do not give a full proof, since it is almost the same as the one given by Rivin and Hodgson [RH93]; the only difference is that the surfaces considered here have genus $g \geq 2$, rather than 0, but this does not appear in the proof.

The main point is that the space of smooth metrics on S with curvature $K < 1$ and contractible closed geodesics of length $L > 2\pi$ is connected. Indeed, given two such metrics h_0 and h_1 , one can take any path connecting them in the space of Riemannian metrics on S , and then "scale up" the middle part, to make sure that the curvature remains small and the closed geodesics remain large (this is also explained in [Sch94, Sch96]).

To prove the first part of the lemma, one first shows that g_0 and g_1 can be approximated by smooth metrics h_0 and h_1 satisfying the curvature and geodesic length conditions. One thus obtains a path $(h_t)_{t \in [0,1]}$ of metrics satisfying the same conditions. One then proves that there is an integer N' such that the metrics in $(h_t)_{t \in [0,1]}$ can be approximated by polyhedral metrics in $\mathcal{M}(S, N)$ in a continuous way.

The second point can be proved in an analogous way, using the fact that the space of smooth metrics on S with curvature $K < 1$ and contractible closed geodesics of length $L > 2\pi$ is simply connected. $\square$

Theorem 4.22. *Let S be a surface of genus $g \geq 2$, and let h be a spherical cone-metric on S , with negative singular curvature at the singular points. Suppose that all contractible closed geodesics of (S, h) have length $L > 2\pi$. Then there is a unique fuchsian polyhedral embedding of (S, h) into H^3 whose third fundamental form is h .*

Note that the uniqueness here is of course up to global isometries of H^3 . Another remark is that the length condition is necessary by lemma 3.11, because the curvature conditions at the vertices imply that the image is convex. By the way, it would be interesting to know whether a similar result also holds with S_1^3 replaced by the anti-de Sitter space.

Proof. As mentioned above, we already know that Φ^C is a local diffeomorphism. Lemma 4.12 shows that Φ^C is proper, so it is a covering of the connected components of $\mathcal{M}^C$ which intersect its image. The first part of lemma 4.21 shows that all of $\mathcal{M}^C$ is in the image, while the second part, along with the fact that $\mathcal{P}^C$ is connected, indicates that each point of $\mathcal{M}$ has a unique inverse image. $\square$

4.4 Fuchsian ideal manifolds

We have already mentioned above that the dihedral angles of ideal polyhedra — and of ideal fuchsian polyhedral embeddings, etc — is the analog of the third fundamental form of finite polyhedra (and of finite fuchsian polyhedral embeddings, etc). Theorem 4.22 should therefore have an analog for ideal fuchsian polyhedral embeddings in terms of dihedral angles. We will prove first the existence part of this statement; the more general result concerning ideal fuchsian polyhedra will be a consequence of other results proved here (specifically, of theorem 8.15) but the existence result given here will be necessary to prove the more general statement. Moreover, we will consider here the case of semi-ideal polyhedral embeddings, which is not covered by theorem 8.15.

Lemma 4.23. *Let Γ be the 1-skeleton of a cellulation of a surface S of genus $g \geq 2$. Let $w : \Gamma_1 \rightarrow (0, \pi)$ be a function on the set Γ_1 of edges of Γ such that:*

1. *for each elementary circuit in Γ , the sum of the values of w is equal to 2π ;*
2. *for each non-elementary circuit, the sum of the values of w is strictly above 2π .*

Then there is a fuchsian ideal embedding of (S, h) into H^3 whose combinatorics is given by Γ , with exterior dihedral angles given by w .

The key point for the remainder of the paper is that, as a consequence, the space of possible dihedral angle assignments is connected — because the space of possible ideal fuchsian polyhedral embeddings is connected.

This lemma is actually a consequence, using property 3.17, of the more general statement below, so we don't prove it separately.

Lemma 4.24. *Let S be a surface of genus $g \geq 2$, and let h be a spherical cone-metric on S , with negative singular curvature at the singular points. Suppose that all contractible closed geodesics of (S, h) have length $L > 2\pi$, except when they bound a hemisphere. Then there is a fuchsian polyhedral embedding of (S, h) into H^3 whose third fundamental form is h .*

Proof. We choose a sequence of spherical cone-metrics $(h_n)_{n \in \mathbb{N}}$ such that:

1. h_n converges to h .
2. for each n , h_n is a spherical cone-metric on S , with negative singular curvature at the singularities.
3. for each n , the contractible closed geodesics of (S, h_n) have length strictly above 2π .

It is not difficult to find such an approximating sequence; one has to decrease slightly the length of some of the edges, and to replace the hemispheres by interiors of convex polygons in S^2 .

Then apply theorem 4.22 to obtain, for each n , a fuchsian finite polyhedral embedding of S in S^3_1 . Finally, lemma 4.12 shows that, after renormalizing this sequence and taking a subsequence, it converges to a semi-ideal fuchsian polyhedral embedding inducing the metric h on S . $\square$

Mathias Rousset [Rou02] recently remarked that the uniqueness part of this statement can be obtained in a rather straightforward way, by using the infinitesimal Pogorelov map to show an infinitesimal rigidity result for semi-ideal fuchsian polyhedra; the uniqueness for semi-ideal polyhedra then follows from studying the map sending a fuchsian polyhedron to its third fundamental form in the neighborhood of semi-ideal polyhedra. He thus obtained that:

Theorem 4.25 (Rousset [Rou02]). *Let S be a surface of genus $g \geq 2$, and let h be a spherical cone-metric on S , with negative singular curvature at the singular points. Suppose that all contractible closed geodesics of (S, h) have length $L > 2\pi$, except when they bound a hemisphere. Then there is a unique fuchsian polyhedral embedding of (S, h) into H^3 whose third fundamental form is h .*

5 Triangulations

This section deals with questions concerning triangulations of a given ideal hyperbolic manifold. Although the existence of a triangulation inducing a given triangulation of the boundary might appear natural at first sight, it is not easy to prove — at least this is not proved here. This is similar to the situation concerning finite volume hyperbolic manifolds, where an ideal triangulation would be helpful but is not known to exist in general; see [PP00]. We will only prove that any ideal hyperbolic manifold M is “almost triangulable” in the sense that it has a finite cover $\overline{M}$ which does admit an ideal triangulation.

Note that alternative approaches could perhaps be followed. One is based on the fact that the main property of the ideal simplices which is used here, namely that the volume is a concave function, remains valid for more general ideal polyhedra (this is a result of [Riv94]). Another (which was pointed out by Francis Bonahon) uses a triangulation which might include some degenerate simplices. Lemma 2.5 does not apply for those simplices, since the volume function is concave but not strictly concave in those cases. It might however be possible to prove that the sum of the volumes of the simplices in a triangulation remains strictly concave, which is basically what one needs in the next sections.

Although those other approaches could presumably lead to shorter proofs, we chose the method described here because, once some technical points are proved, it gives rather simple picture of what goes on; and also because the method it contains might be useful in other settings.

5.1 Triangulations, cellulations

We first give more details about what we call a triangulation here.

Definition 5.1. *Let M be an ideal hyperbolic manifold. A **cellulation** C of M is a finite family $C_1, C_2, \dots, C_n$ of non-degenerate, closed, convex, ideal polyhedra isometrically embedded in M , such that:*

1. *for $i \neq j$, the interiors of C_i and C_j are disjoint;*
2. *the union of the C_i is all of M ;*
3. *for $i \neq j$, if $C_i \cap C_j \neq \emptyset$, then it is a face of both C_i and C_j .*

The C_i are the “cells” of the cellulation C .

The third condition excludes some “bad” configuration, like the one depicted in figure 1, where two simplices have an intersection which is not a face in any of them.




Figure 1: This is forbidden in our cellulations

Definition 5.2. *A **triangulation** of M is a cellulation whose cells are all simplices.*

Definition 5.3. *An ideal hyperbolic manifold M is **triangulable** if it admits a triangulation. It is **almost triangulable** if it has a finite cover which is triangulable.*

The main goal of this section will be to prove the:

Lemma 5.4. *Any ideal hyperbolic manifold is almost triangulable.*

The proof will proceed in several steps. The first point is the:

Proposition 5.5. *Any ideal hyperbolic manifold admits a cellulation.*

Proof. It is done along the ideas of Epstein and Penner [EP88]; the situation here is simpler since the action of $\pi_1 M$ on $S^2 \setminus \Lambda$ is discrete.

Let M be an ideal hyperbolic manifold. Then M is isometric to the convex hull of a set $\{x_1, \dots, x_N\}$ of points in $\partial_\infty E(M)$, where $E(M)$ is the unique convex co-compact hyperbolic manifold in which M admits an isometric embedding which is surjective on the π_1 .

For each $i \in \{1, 2, \dots, N\}$, choose a “small” horoball $b_i \subset E(M)$ with ideal point x_i ; we suppose that the b_i are small enough to be disjoint. Let B_i be the lift of b_i to H^3 , which is $\pi_1 M$ -invariant collection of disjoint horoballs in H^3 .

Now we want to use this action of $\pi_1 M$ on the horoballs to produce a cellulation of M . This is done in [EP88] by considering the action of $\pi_1 M$ on the light cone in Minkowski 4-space, which contains H^3 as a quadric. We will use here a similar, slightly more complicated but maybe a little more geometric, approach. It is based more explicitly on the action of $\pi_1 M$ on the space of horospheres, with an explicit model from [Sch02b].

We consider the projective model of H^4 and half of S_1^4 , the de Sitter space of dimension 4, corresponding to the 3-dimensional models already described above. It can be obtained as follows. H^4 and S_1^4 are both isometric to quadrics in Minkowski 5-space, with the induced metric:

$$H^4 \simeq \{x \in \mathbf{R}_1^5 | \langle x, x \rangle = -1 \wedge x_0 > 0\} ,$$

$$S_1^4 \simeq \{x \in \mathbf{R}_1^5 | \langle x, x \rangle = 1\} .$$

Let P_0 be the affine hyperplane of equation $x_0 = 1$ in $\mathbf{R}_1^5$; consider the map sending a point $x \in H^4$ (resp. $x \in S_1^4$ with $x_0 > 0$) to the intersection with P_0 of the line going through x and 0. It is not difficult to check that ϕ is projective; it maps H^4 to the interior of the radius one ball, and the part of S_1^4 standing on one side of a space-like hyperplane to its exterior. We now consider this model only, with P_0 identified with $\mathbf{R}^4$.

Using the classical Poincaré model of H^3 , we can map conformally H^3 to the interior of a geodesic ball B_0 in S^3 , e.g. to the hemisphere:

$$S_+^3 := \{x \in \mathbf{R}^4 | \langle x, x \rangle = 1 \wedge x_1 \geq 0\} .$$

Horospheres in H^3 are then mapped to spheres in S^3 which are interior to S_+^3 and tangent to its boundary. Those spheres are the boundaries at infinity of the totally geodesic 3-planes in H^4 which are asymptotic to a given 3-plane H_1 , with $\partial_\infty H_1 = \partial S_+^3$. Their dual points in S_1^4 (using the hyperbolic-de Sitter duality, see subsection 3.4) form the vertical cylinder C_0^+ which is tangent to S^3 along ∂S_+^3 . In de Sitter terms, C_0^+ is the future (or past, depending of the orientation) light cone of a point H_1^* which is at infinity in the projective model of S_1^4 which we use.

By construction, the action of $\pi_1 M$ on H^3 extends to a conformal action on S^3 , and thus to a projective action on $\mathbf{R}^4$. This action leaves invariant S_+^3 , and thus also H_1 and H_1^* , and therefore also C_0^+ . For each $i \in \{1, \dots, N\}$, the horoballs in B_i corresponds to the points of an orbit O_i of the action of $\pi_1 M$ on C_0^+ . Since the horoballs in B_i are disjoint, it is easy to see that O_i is discrete, with no accumulation point outside ∂S_+^3 .

To finish the proof, we proceed almost as in [EP88]; consider the convex hull (in $\mathbf{R}^4$) of $O_1 \cup \dots \cup O_N \cup H_1$, which by the property just pointed out is locally finite outside $\partial S_+^3 = \partial H_1$, with faces which are polyhedra with a finite number of edges and vertices. Then take the “projection” of this polyhedral surface on $H^3 \subset S^3$ in the vertical direction to obtain the required cellulation. By construction it is invariant under the action of $\pi_1 M$ on H^3 .

The cellulation of M obtained in this way has a finite number of cells. Otherwise, there would exist an edge e meeting an infinite number of fundamental domains of the action of $\pi_1 M$ on $\tilde{M}$. So e would connect a vertex v of $\tilde{M}$ to the limit set Λ of the action of $\pi_1 M$ on H^3 . Now note that, by construction, M is covered by a finite set of (non-disjoint) ideal simplices $T_1, \dots, T_N$. So there would exist $i \in \{1, \dots, N\}$ such that e intersects γT_i for an infinite set S of elements γ of $\pi_1 M$. Going back to the projective model




Figure 2: The space of horospheres as a cone

used above, for each $\gamma \in S$, the segment γe should intersect T_i — and it would also be in the boundary of the convex hull constructed above.

Now γe goes from γv to a point $\gamma v'$ of Λ ; as $\gamma \rightarrow \infty$, the vertical coordinates of both γv and $\gamma v'$ go to zero, so that the whole segment γe goes to the horizontal hyperplane containing 0. Therefore, γe lies "below" T_i , so that the intersection $\gamma e \cap T_i$ can not be in the boundary of the convex hull. So such an e can not exist, and the cellulation obtained has a finite number of cells. $\square$

5.2 From a cellulation to a triangulation

We now consider a cellulation C of an ideal hyperbolic manifold M , and call F_0, F_1, F_2 and F_3 the sets of its faces of dimension 0, 1, 2 and 3 respectively. We also consider the universal cover $\tilde{M}$ of M ; C lifts to a cellulation $\tilde{C}$ of $\tilde{M}$, and we call $\tilde{F}_0, \tilde{F}_1, \tilde{F}_2$ and $\tilde{F}_3$ the sets of its faces of the 4 possible dimensions.

From now on, and until section 8, we will consider a fixed triangulation Σ on M .

Definition 5.6. Let C be a cellulation of an ideal hyperbolic manifold M . C is **large** if, for any cell $\sigma \in \tilde{F}_3$, any vertex s of σ and any $\gamma \in \pi_1(M)$, if γs is again a vertex of σ , then $\gamma = 1$.

Proposition 5.7. For any ideal hyperbolic manifold M and any cellulation C of M , M has a finite cover $\overline{M}$ such that C lifts to a large cellulation.

Definition 5.8. Let C be a cellulation of an ideal hyperbolic manifold M . A **polarization** of C is a map $\rho : \tilde{F}_3 \rightarrow \tilde{F}_0$ which is equivariant under the action of $\pi_1 M$ and such that:

1. for any $\sigma \in \tilde{F}_3$, $\rho(\sigma)$ is a vertex of σ ;
2. if $\sigma \in \tilde{F}_3$ and $\sigma' \in \tilde{F}_3$ are adjacent (i.e. $\sigma \cap \sigma' \in \tilde{F}_2$), and if $\rho(\sigma)$ and $\rho(\sigma')$ are both vertices of $\sigma \cap \sigma'$, then $\rho(\sigma) = \rho(\sigma')$.

Let's pause to remark that, although a $(\pi_1 M)$ -equivariant map from $\tilde{F}_3$ to $\tilde{F}_0$ induces a map from F_3 to F_0 , the equivariant map contains much more information. Indeed this is already apparent in the simple case where one considers the manifold S^1 , triangulated with only one edge and one vertex. Its universal cover is $\mathbf{R}$, triangulated with vertices at the integers. There are two $\mathbf{Z}$ -equivariant maps sending a segment $[k, k+1]$ to one of its endpoints: $\rho_1 : [k, k+1] \mapsto k$, and $\rho_2 : [k, k+1] \mapsto k+1$. There is however only one map from F_3 to F_0 , since F_0 has only one element.

Proposition 5.9. Any large cellulation C of an ideal hyperbolic manifold M admits a polarization.

Proposition 5.10. Any polarized cellulation of an ideal hyperbolic manifold M can be subdivided to obtain a triangulation. If the cellulation is large, so is the triangulation obtained.

The proof of lemma 5.4 clearly follows from the three propositions above so it remains only to prove them.

Proof of proposition 5.7. Let $\sigma \in \tilde{F}_3$. First note that, if $\gamma \in \pi_1 M$ fixes the vertices of σ , then $\gamma = e$; indeed, no non-identity element of $\text{Isom}(H^3)$ has more than 2 fixed points at infinity. Therefore, the elements of $\pi_1 M$ leaving σ invariant are determined by their actions on its vertices, so that there is a finite subset E_σ of $\pi_1 M$ of elements leaving σ invariant.

Now C has a finite number of cells, which we can call $\sigma_1, \dots, \sigma_N$. For each i , let $\bar{\sigma}_i$ be a cell of $\tilde{C}$ whose projection on M is σ_i . Then $E := \cup_i E_{\bar{\sigma}_i}$ is finite, where E is the set of elements of $\pi_1 M$ leaving one of the $\bar{\sigma}_i$ invariant. Since there is a finite number of cells $\bar{\sigma}'$ sharing a vertex with some given cell $\bar{\sigma}$, the set $F_{\bar{\sigma}}$ of elements γ of $\pi_1 M$ such that $\bar{\sigma}$ shares a vertex with $\gamma\bar{\sigma}$ is finite. Therefore, the set $F := \cup_i F_{\bar{\sigma}_i}$ is finite.

$\pi_1 M$ is linear, and therefore residually finite (see e.g. [LS01], chapter III, 7.11). Thus there exists a normal subgroup Γ of $\pi_1 M$ of finite index, such that $\Gamma \cap F = \{1\}$. The corresponding finite cover $\bar{M}$ of M has the required property. $\square$

Proof of proposition 5.9. We will construct the required polarization ρ as the endpoint of a sequence of partially defined equivariant functions $\rho_i : \tilde{F}_3 \rightarrow \tilde{F}_0$ (that is, functions defined on a subset of $\tilde{F}_3$ only).

First choose $s_0 \in \tilde{F}_0$; since C is large, for any $\sigma \in \tilde{F}_3$, at most one of the vertices of σ is in $(\pi_1 M)s_0 \subset \tilde{F}_0$. Define ρ_0 on a cell $\sigma \in \tilde{F}_3$ as follows:

- if there exists $\gamma \in \pi_1 M$ such that γs_0 is a vertex of σ , then set $\rho_0(\sigma) := \gamma s_0$;
- otherwise, leave ρ_0 undefined at σ .

It is clear that this partially defined map is equivariant.

Now choose $s_1 \in \tilde{F}_0$ such that some $\sigma \in \tilde{F}_3$ on which ρ_0 is not defined has s_1 as a vertex, and define ρ_1 as follows:

- if there exists $\gamma \in \pi_1 M$ such that γs_1 is a vertex of σ , then $\rho_1(\sigma) := \gamma s_1$;
- otherwise, $\rho_1(\sigma) := \rho_0(\sigma)$.

The second case includes the possibility that ρ_0 is undefined at σ , then ρ_1 remains undefined at σ .

Then repeat this construction with s_2 to obtain a map ρ_2 , etc. The number of cells of F_3 on which ρ_i is not defined decreases by at least one unit at each step, and C has a finite number of cells, so after a finite number of steps we obtain an equivariant map $\rho := \rho_N : \tilde{F}_3 \rightarrow \tilde{F}_0$ which is everywhere defined.

We now want to prove that ρ is a polarization. It is clear by construction that, for any $\sigma \in \tilde{F}_3$, $\rho(\sigma)$ is a vertex of σ . Let $\sigma' \in \tilde{F}_3$ be another cell, such that $\sigma \cap \sigma' \in \tilde{F}_2$. Let:

$$i_0 := \max\{i \in \{1, \dots, N\} \mid \exists \gamma \in \pi_1 M, \gamma s_i \text{ is a vertex of } \sigma\} ,$$

$$j_0 := \max\{j \in \{1, \dots, N\} \mid \exists \gamma' \in \pi_1 M, \gamma' s_j \text{ is a vertex of } \sigma'\} .$$

We consider two cases:

1. γs_{i_0} is a vertex of σ but not of σ' ; then $\rho(\sigma) = \gamma s_{i_0}$ is not a vertex of σ' , and condition (2) of definition 5.8 is satisfied. The same applies if $\gamma' s_{j_0}$ is not a vertex of σ .
2. γs_{i_0} and $\gamma' s_{j_0}$ are both vertices of both σ and σ' . But then, by definition of i_0 and j_0 , $i_0 = j_0$ and $\rho(\sigma) = \gamma s_{i_0} = \gamma' s_{j_0} = \rho(\sigma')$, so that condition (2) of definition 5.8 again applies.

$\square$

Proof of proposition 5.10. Let C be a cellulation of M , with a polarization ρ . We first built a $\pi_1 M$ -invariant triangulation of $\tilde{F}_2$ as follows.

Let $f \in \tilde{F}_2$ be such that some vertex s of f is $\rho(\sigma)$, where σ is one of the cells bounded by f . Then s is the unique vertex of f with this property, because of condition (2) of definition 5.8. Define a triangulation of f by adding the edges going from s to all the other vertices of f . Repeat this for all the 2-faces with this property.

Then subdivide all the remaining non-triangular 2-faces of F_2 , so as to obtain an equivariant triangulation of $\tilde{F}_2$.

Finally, define a triangulation of M by subdividing each cell σ of $\tilde{F}_3$ by adding triangles containing $\rho(\sigma)$ and any edge of σ not containing $\rho(\sigma)$. It is clear that:

1. this defines an equivariant decomposition of $\tilde{M}$ into simplices, which is obtained by subdividing each cell into simplices, and thus a decomposition of M into a finite number of simplices with disjoint interior;
2. the simplices are non-degenerate;
3. if $\sigma \in \tilde{F}_3$ and $\sigma' \in \tilde{F}_3$ are two adjacent simplices, then $\sigma \cap \sigma'$ is a face in both of them, because it has to be one of the triangles of the triangulation of $\tilde{F}_2$ obtained above.

The definition then directly shows that the triangulation obtained in this manner from a large cellulation is itself large. $\square$

5.3 Some elementary combinatorics

We now fix a triangulation Σ of M , with which we will stick until section 8. We call f the number of its 3-simplices, t the number of its 2-faces, e the number of its edges, e_i and e_b the number of interior and boundary edges respectively, and v the number of vertices. We will need later on the following easy consequence of the Euler formula.

Lemma 5.11. $2f = 2e_i + e_b - v$.

Proof. Consider the closed triangulated manifold obtained by gluing two copies of (M, Σ) along their boundary by the identity map. This triangulated manifold has $\bar{f} := 2f$ simplices, $\bar{t}$ 2-faces, $\bar{e} := 2e_b + e_i$ edges, and $\bar{v} := v$ vertices.

Since the Euler characteristic is 0 in odd dimensions:

$$\bar{f} - \bar{t} + \bar{e} - \bar{v} = 0 .$$

Moreover, each 2-face bounds two simplices, and each simplex has 4 faces, so that:

$$\bar{t} = 2\bar{f} .$$

Therefore:

$$-\bar{f} + \bar{e} - \bar{v} = 0 ,$$

and the result follows. $\square$

6 Hyperbolic structures on triangulated manifolds

This section contains the definitions and basic properties concerning some simple notions of singular hyperbolic structures. The idea is to construct such structures by gluing ideal simplices, and then to show that the set of those structures with some constraints actually contains a smooth hyperbolic metric. This will be done in the next section using a variational argument.

The ideas used here were mostly developed previously for ideal polyhedra in H^3 . Their history is interesting. The first results were obtained by Andreev [And71], and then developed by Thurston [Thu97]. Colin de Verdière [CdV91] then noted that the results could be recovered using a variational approach, while Brägger [Brä92] identified the functional as the volume. Rivin [Riv94] then further developed the theory.

6.1 Sheared hyperbolic structures

We consider here a triangulation Σ of M .

Definition 6.1. A **sheared hyperbolic structure** on (M, Σ) is the choice, for each simplex S in Σ , of a diffeomorphism from S to an ideal simplex. We denote by $\mathcal{H}_{sh}$ the set of hyperbolic structures on (M, Σ) .

This vocabulary is justified by the elementary remark that, since all ideal triangles in H^2 are isometric, there is a unique way of gluing the hyperbolic simplices which are given by a sheared hyperbolic structure along their common faces. One then obtains a hyperbolic metric on the complement of the interior edges of Σ . But this metric does not extend over the edges; rather, the model of what happens along an edge is obtained by taking the quotient of the universal cover of H^3 minus a geodesic g by the group generated by the composition of a rotation of angle θ around g and a translation of length δ along g . We then call θ the **angle** around the edge, and δ the **shear** along the edge.

Those hyperbolic structures will be considered up to diffeomorphisms acting on the simplices; therefore, the choice of a sheared hyperbolic structure is equivalent to the choice of the three dihedral angles of each simplex, subject to the condition that their sum is π . So $\mathcal{H}$ can be identified with the product of f 2-simplices, and it has a natural affine structure.

Definition 6.2. A **sheared hyperbolic structure** on (M, Σ) is a **singular hyperbolic structure** if the shear of all the interior edges vanishes. It is **smooth** if, in addition, the singular angle of all interior edges is 2π . The set of singular hyperbolic structures is denoted by $\mathcal{H}_{si}$, the set of smooth hyperbolic structures by $\mathcal{H}_{sm}$.

In other words, a singular hyperbolic structure on (M, Σ) defines a hyperbolic cone-manifold structure on M , which is singular on the edges of Σ . A smooth hyperbolic structure defines a hyperbolic metric on M . In both cases the boundary is piecewise totally geodesic.

The shear at an interior edge of Σ can be understood in the following elementary way.

Definition 6.3. Let s be an ideal simplex, and let e be an edge of s . Choose an orientation of e , and let f_1 and f_2 be the faces of s containing e , in the order defined by the orientation of e . Let x_1 and x_2 be the orthogonal projections on e of the vertices of f_1 and f_2 respectively which are not in e . The **shear** of s at e is the oriented distance between x_1 and x_2 .

Note that the shear of a simplex at an edge is clearly independent of the orientation chosen.

Remark 6.4. Let $h \in \mathcal{H}_{sh}$. The shear of h at an interior edge e is the sum, over the simplices containing e , of their shears at e .

6.2 Angles

For each edge e of an ideal triangulation of a sheared hyperbolic structure, we define the **angle** at e to be the sum of the dihedral angles at that edge of the simplices containing it – this applies to interior as well as to boundary edges. For boundary edges this will also be called the **interior dihedral angle**, and the **exterior dihedral angle** is π minus the interior dihedral angle. For interior edges, the **excess angle** is the angle minus 2π , and the **singular curvature** around the corresponding edge is minus the excess angle.

Note that, for $h \in \mathcal{H}_{sh}$, the sum of the exterior angles at a vertex is equal to 2π plus the sum of the excess angles at the interior angles. This is checked by applying the Gauss-Bonnet formula to the link of the vertex, which is piecewise Euclidean manifold.

Definition 6.5. Let $\theta : \Sigma_1 \rightarrow \mathbf{R}_+$ be an assignment of “angles” to the edges of Σ . We will say that θ is “ideal” if:

- the angles assigned to boundary edges are in $(0, \pi)$;
- the angles assigned to interior edges are in $(0, 2\pi)$;

- at each vertex, the sum of the exterior dihedral angles of the boundary edges equals 2π plus the sum of the angle excess at the interior angles.

The set of ideal angle assignments is denoted by Θ . For each $\theta \in \Theta$, we denote by $\mathcal{H}_{sh}(\theta)$ the set of sheared hyperbolic structures on Σ such that the angles associated to each edge is given by θ ; then $\mathcal{H}_{si}(\theta) := \mathcal{H}_{sh}(\theta) \cap \mathcal{H}_{si}$, and $\mathcal{H}_{sm}(\theta) := \mathcal{H}_{sh}(\theta) \cap \mathcal{H}_{sm}$.

Lemma 6.6. *If the triangulation Σ is large, then Θ corresponds to the interior of a polytope of dimension $e - v$ in $\mathbf{R}^e$.*

Proof. The only point is to prove that the constraints on the vertices are linearly independent. So let C be linear combination of those constraints which is zero. In other terms, C is a function $C : \Sigma_0 \rightarrow \mathbf{R}$ such that, for each oriented edge e , $C(e_-) + C(e_+) = 0$. Moreover we supposed that Σ is large, so it is quite obvious that the values of C at the vertices of a triangular face of Σ have to be 0, and therefore that $C = 0$. $\square$

Definition 6.7. *We say that an ideal angle assignment is **smooth** if the angle assigned to each interior edge is 2π . The set of those angle assignments is denoted by Θ_{sm} .*

7 First order variation of the volume

The main goal of this section is to use elementary properties of the volume — seen as a functional on the space $\mathcal{H}_{sh}$ of sheared hyperbolic structures — to prove the existence of singular hyperbolic structures with given angles on the interior and boundary edges. In the next section, the results and some similar arguments will be used to understand the smooth hyperbolic structures with given dihedral angles at the boundary edges.

7.1 Definitions and first properties

We consider here again an ideal hyperbolic manifold, along with a triangulation Σ , and an angle assignment $\theta \in \Theta$. We first define the volume of a singular hyperbolic structure in the most obvious way.

Definition 7.1. *For $h \in \mathcal{H}_{sh}$, the **volume** of h , $V(h)$, is the sum of the hyperbolic volumes of the simplices of Σ .*

As an immediate consequence of lemma 2.5, we find that:

Lemma 7.2. *V is a concave function on $\mathcal{H}_{sh}$ (with the affine structure coming from the parametrization by the dihedral angles of the simplices).*

We now need to understand how the volume varies when one deforms a sheared hyperbolic structure. Unfortunately the Schläfli formula (1) does not apply directly to sheared hyperbolic structures, and does not even make sense in this case; indeed there is no way to choose a horosphere centered at a given vertex, since the holonomy around an edge e would act on it by translation along the edge, with a translation distance equal to the shear at e .

To understand this point better, we fix a sheared hyperbolic structure $h \in \mathcal{H}_{sh}(\theta)$, and a first order variation $\dot{h} \in T_h \mathcal{H}_{sh}(\theta)$. Suppose given an ideal triangulation of h . $\dot{h}$ determines a first order variation of the dihedral angles of the simplices, to which the Schläfli formula (1) can be applied, once a horosphere around each vertex is chosen for each simplex. To get a better understanding of the first order variation of the volume, we can choose, for each vertex v , a horosphere centered on v for each simplex containing v . We call this collection of horospheres a **choice of horospheres** at v .

Definition 7.3. *Let e be an edge of Σ , with vertices e_- and e_+ . Let H_- and H_+ be choices of horosphere at e_- and e_+ respectively. The **volume variation at e** associated to $\dot{h}$ is:*

$$\sum_i L_i \dot{\alpha}_i,$$

where the sum is over the simplices containing e , L_i is the oriented length of the part of e which is between the horospheres in H_- and H_+ , and $\dot{\alpha}_i$ is the first-order variation of the dihedral angle at e .

7.2 Volume differential and shears

Of course the point is that, once horospheres are chosen, the volume variation at e can be seen as the contribution coming from e to the first order variation of the volume; indeed, the Schläfli formula (1) indicates that:

Remark 7.4. *If a horosphere choice is given for all vertices of Σ , the first order variation of the volume under the deformation $\dot{h}$ is the sum of the volume variations at the edges.*

Definition 7.5. *Let T be a 2-face of the triangulation Σ , and let v be a vertex contained in Σ . A choice of horospheres at v is **coherent** at T if the horospheres on both simplices containing T have the same intersection with T .*

The point is that, as explained above, if h has a non-zero shear at an edge e containing v , then it is not possible to find a horosphere choice at v which is coherent on all 2-faces containing e .

Proposition 7.6. *Let e be an oriented interior edge of Σ , with endpoints e_- and e_+ . Let $f_1, \dots, f_N$ be the 2-faces containing e , oriented in cyclic order. Choose $i, j \in \{1, \dots, N\}$. Let H_- be a choice of horospheres at e_- around e which is coherent except at f_i , and let H_+ be a choice of horospheres at e_+ around e which coherent except at f_j . Let $\dot{h}$ be an infinitesimal variation of h which does not change the total angle around the interior edges. The variation of the volume at e associated to $\dot{h}$ is:*

$$\dot{V}_e = s_e \sum_{k=j}^{i-1} \dot{\alpha}_k ,$$

where s_e is the shear of h at e , and α_k is the angle between f_k and f_{k+1} .

Proof. The definition of H_- and H_+ shows that there are two numbers $L, L' \in \mathbf{R}$ such that:

- for each $k \in \{i, i+1, \dots, j-1\}$, the distance along e between the horospheres of H_- and H_+ in the simplex having faces f_k and f_{k+1} is L .
- for each $k \in \{j, j+1, \dots, i-1\}$, the same distance is L' .

Then $L' - L = s_e$, again by definition of H_- , H_+ and s_e . So, by definition 7.3:

$$\dot{V}_e = \sum_{k=i}^{j-1} L \dot{\alpha}_k + \sum_{k=j}^{i-1} L' \dot{\alpha}_k = \sum_{k=j}^{i-1} L \dot{\alpha}_k + \sum_{k=j}^{i-1} (L + s_e) \dot{\alpha}_k .$$

But the total angle around e remains constant in the variation $\dot{h}$, which means that:

$$\sum_k \dot{\alpha}_k = 0 .$$

Subtracting L times this equation to the previous one leads to the statement. $\square$

As a consequence we see that the first-order variation of the volume, under a deformation which does not change the total angle around the interior angles or the dihedral angles, has a remarkably simple form, more precisely it can be expressed only in terms of the shears at the interior edges.

Lemma 7.7. *$dV(\dot{h})$, as a linear form on $\mathcal{H}_{sh}(\theta)$, depends only on the shear at the interior edges of Σ .*

Proof. This is an immediate consequence of the previous proposition, and of the fact that the dihedral angle at the exterior edges is constant. $\square$

This means in particular that the volume — seen as a function on $\mathcal{H}_{sh}(\theta)$ — is critical when the shear vanishes at all interior edges. We will see below that the converse is true too. This will use a special type of deformations, defined as follows.

Definition 7.8. Let $h \in \mathcal{H}_{si}$, and let $e \in \Sigma_1$ be an oriented interior edge of Σ . A **local** deformation $\dot{h}_0$ of h at e is determined as follows. Choose a face f containing e . Then let $s_0, s_1, s_2, \dots, s_N = s_0$ be the simplices of Σ containing e , in cyclic order. Let a_i^+ and b_i^+ be the edges of s_i containing e_+ (other than e), and a_i^- and b_i^- the edges of s_i containing e_- (again other than e), ordered so that $a_i^+ = b_{i+1}^+$ and that $a_i^- = b_{i+1}^-$. Then, in the deformation $\dot{h}_0$:

- the angles of the simplices s_i at e do not change;
- the angles of s_i at a_i^+ and b_i^- vary at speed 1;
- the angles of s_i at a_i^- and at b_i^+ vary at speed -1 .




Figure 3: Local deformations

It is a simple matter to check the following properties of local deformations:

Remark 7.9. The above definition indeed defines a deformation of h in $\mathcal{H}_{sh}$. Moreover, it does not change the total angles at the edges, i.e. if $h \in \mathcal{H}_{sh}(\theta)$, then $\dot{h} \in T_h \mathcal{H}_{sh}(\theta)$.

The local deformations can then be used to prove that, among the sheared hyperbolic structures, those which have zero shear are exactly the critical points of the volume, seen as a functional on the space of sheared hyperbolic structures having a given total angles on all edges.

Lemma 7.10. Let $\theta \in \Theta$, and let $h \in \mathcal{H}_{sh}(\theta)$; then the restriction of dV to $T_h \mathcal{H}_{sh}(\theta)$ is zero if and only if the shear of h at all interior edges is zero, that is, if and only if $h \in \mathcal{H}_{si}$.

Proof. Lemma 7.7 shows that the volume — as a functional on $\mathcal{H}_{sh}(\theta)$ — is critical when the shear is zero at all interior edges. Remark 7.9 shows that the local deformations are tangent to $\mathcal{H}_{sh}(\theta)$.

Suppose that the shear at an edge e is non-zero, and consider the simplices containing e , as in figure 3. It is possible to choose horospheres at the vertices of those simplices such that:

- at e_+ and e_- , the choices are coherent at the triangles (e, a_i^+, a_i^-) , except for $i = 0$;
- at e_+ and e_- , the distance between the two horospheres on each side of the triangle (e, a_0^+, a_0^-) is equal to the shear at e ;
- for each $i \neq 0$, the two horospheres at the vertex $a_i^- \cap a_i^+$ are coherent.

Consider a local deformation at e . The only contributions to the variation of the volume comes from the a_i^- and the a_i^+ , since they are the only edges where the angles vary. Moreover they add up to zero except at a_0^- and a_0^+ , because of the choices of horospheres described above (they are coherent on both sides of the other edges). Finally, the contributions from a_0^- and a_0^+ are non-zero and of the same sign.

So proposition 7.6 shows that the shears have to be zero at all interior edges at critical points of V . $\square$

7.3 Consequences

The results of the previous subsection now easily lead to interesting results on singular hyperbolic structures (i.e. those without shears).

Corollary 7.11. *Let $\theta_0 \in \Theta$ and $g_0 \in \mathcal{H}_{si}(\theta_0)$. There exists a neighborhood U of θ_0 in Θ and a neighborhood V of g_0 in $\mathcal{H}_{sm}$ such that, for any $\theta \in U$, there exists a unique $g \in \mathcal{H}_{si}(\theta) \cap V$.*

Proof. By corollary 2.6, V is a strictly concave function on $\mathcal{H}_{sh}$. Moreover, the restriction of V to $\mathcal{H}_{sh}(\theta_0)$ has a critical point at g_0 . Therefore, for θ close enough to θ_0 , V has a unique critical point of V on $\mathcal{H}_{sh}(\theta)$ close to g_0 ; by lemma 7.10 it is a point in $\mathcal{H}_{si}(\theta)$. $\square$

Corollary 7.12. *For each $\theta \in \Theta$ such that $\mathcal{H}_{sh}(\theta)$ is non-empty, $\mathcal{H}_{sh}(\theta)$ is a submanifold of $\mathcal{H}_{sh}$, with dimension equal to the number e_i of interior edges of the triangulation.*

Proof. The space of ideal simplices in H^3 has dimension 2, so that the dimension of $\mathcal{H}_{sh}$ is equal to twice the number of simplices in Σ . Thus, by lemma 5.11, $\dim \mathcal{H}_{sh} = 2e_i + e_b - v$, where e_b and v are the number of boundary edges and of vertices of Σ respectively.

Specifying the dihedral angles and the excess angle at the interior edges adds $e_i + e_b$ constraints; they are not linearly independent, however, since they satisfy at least one linear condition for each vertex. The loss of dimensions due to those constraints is therefore at most $e_i + e_b - v$, so that, for $\theta \in \Theta$, $\dim \mathcal{H}_{sh}(\theta) \geq e_i$.

But lemma 7.7 indicates that dV depends only on the e_i shears at the interior edges, while the strict concavity of V shows that the Jacobian of dV is non-degenerate. Therefore, its restriction to $\mathcal{H}_{sm}(\theta)$ is also non-degenerate, so that $\dim \mathcal{H}_{sh}(\theta) \leq e_i$, and the result follows. $\square$

Finally, using the non-degeneracy of the volume functional also leads to an infinitesimal rigidity result.

Corollary 7.13. *Let $h \in \mathcal{H}_{si}$, and let $\dot{h} \in T_h \mathcal{H}_{si}$ be a first-order deformation of h . Then $\dot{h}$ corresponds to a non-zero first-order deformation of either the dihedral angles of the boundary edges, or the excess angle at the interior edges. $\mathcal{H}_{si}$ is, in the neighborhood of h , a manifold of dimension $e_i + e_b - v$.*

Proof. This is a direct consequence of corollary 7.12, which shows that $\mathcal{H}_{sh}$ is foliated by the $\mathcal{H}_{sh}(\theta)$, which are submanifolds of dimension e_i . We already know that $\mathcal{H}_{sh}$ is a manifold of dimension $2f$, and, by lemma 5.11, $2f = 2e_i + e_b - v$, where e_b is the number of boundary edges and v is the number of vertices. But the dimension of the space Θ of possible angle assignments is at most $e_b + e_i - v$, since the dihedral angles at the boundary edges satisfy one linear condition for each vertex. Thus this dimension has to be exactly $e_b + e_i - v$, and each non-trivial deformation in $\mathcal{H}_{si}$ — which is transverse to the foliation of $\mathcal{H}_{sh}$ by the $\mathcal{H}_{sh}(\theta)$ — induces a non-trivial deformation of some dihedral angle on the boundary or of some total angle on an interior edge. $\square$

Finally, an elementary dimension-counting argument, along with the previous corollary, shows that any infinitesimal deformation of the dihedral angles on the boundary — and of the angle excess around the interior edges — can be realized by some $\bar{h} \in T_h \mathcal{H}_{si}$, if it satisfies the condition that, at each vertex, the sum of the exterior dihedral angles remains equal to 2π plus the sum of the angle excess at the interior edges. Restricting to the case of smooth hyperbolic structures, this can be formulated as follows. We call **admissible** an infinitesimal deformation of the dihedral angles such that the sum of the exterior angles of the edges at any vertex remains 2π .

Lemma 7.14. *Let (M, h) be an ideal hyperbolic manifold. For any admissible infinitesimal variation $\dot{\theta}$ of the dihedral angles on the boundary, there exists a (unique) infinitesimal deformation of h in $\mathcal{H}_{si}$ which induces the variation $\dot{\theta}$.*

Proof. By lemma 5.4, there exists a finite cover $\bar{M}$ of M so that the hyperbolic structure $\bar{h}$ lifted to $\bar{M}$ from h admits an ideal triangulation, say Σ . Let $\bar{\theta}$ be the dihedral angles of $\bar{h}$. The previous corollary shows that there is no non-trivial deformation of $\bar{h}$ which does not change the dihedral angles, and that the dimension of $\mathcal{H}_{sm}$ in the neighborhood of $\bar{h}$ is $e_b - v$. Thus, for each admissible infinitesimal deformation $\dot{\bar{\theta}}$ of the dihedral angles of $\bar{h}$, there is a unique infinitesimal deformation $\dot{\bar{h}}$ of $\bar{h}$ inducing $\dot{\bar{\theta}}$.

If now $\dot{\theta}$ is an infinitesimal deformation of θ , it lifts to an infinitesimal deformation $\dot{\bar{\theta}}$ of the dihedral angles of $\bar{h}$, which is induced by a unique admissible infinitesimal deformation $\dot{\bar{h}}$ of $\bar{h}$. Since $\dot{\bar{\theta}}$ is equivariant, $\dot{\bar{h}}$ is obviously equivariant (by uniqueness) and therefore defines an admissible infinitesimal deformation $\dot{h}$ of h inducing $\dot{\theta}$, which is unique since $\dot{h}$ is. $\square$

Note that it is not necessary to suppose that the hyperbolic ideal manifold which we consider has a boundary which is triangulated, i.e. this lemma is also valid when some of the faces have more than 3 edges. In this case one can add edges to those faces, so as to obtain triangulation, and then choose the first-order variation of the dihedral angles at those new edges. In particular it is possible to choose first-order variations which "lose" the convexity of the boundary.

8 Dihedral angles

We consider here an ideal hyperbolic manifold M . The first point will be a compactness result for sequences of ideal structures on M with a given triangulation of the boundary, when the dihedral angles converge. We will then consider a fixed triangulation of M , then see what happens when one considers only a fixed cellulation σ of the boundary, and finally how the affine structures corresponding to different triangulations of the boundary can be glued together.

8.1 Compactness of ideal manifolds

We consider a fixed triangulation σ of ∂M ; we will give in this subsection a compactness lemma for ideal structures with boundary combinatorics corresponding to σ , then use it in the next sub-sections to obtain a description of the possible dihedral angles.

Lemma 8.1. *Let (h_i) be a sequence of ideal structures on M with boundary combinatorics given by σ . Suppose that the dihedral angles of the boundary vertices for the h_i converge as $i \rightarrow \infty$ to limits in $(0, \pi)$. Then:*

- *either there is a non-elementary circuit in σ on which the sum of the limit dihedral angles is 2π ;*
- *or there exists a sub-sequence of (h_i) which converges to an ideal structure on M .*

The intuitive idea behind this lemma is simple. Consider the induced metrics on one of the boundary components. For each n , it is a complete hyperbolic metric of finite area. So, by a classical result in Teichmüller theory, there is a subsequence which either converges to a hyperbolic metric of finite area, or has a closed geodesic whose length goes to 0. In the second case, the short geodesic will — by a simple convexity argument — correspond to a non-elementary circuit on which the sum of the exterior dihedral angles goes to 2π . If the first case happens for all boundary components, then both the induced metric and the dihedral angles converge, so that the universal cover in H^3 of each boundary component converges, and thus the sequence of metrics on M has to converge.

The "formal" proof, however, is a little more complicated since one has to keep track of the relationship between the induced metrics and the triangulation; it will use a couple of definitions and propositions. First there is a natural notion of collapsing of vertices for the sequence (h_i) . We consider the triangulation σ of ∂M , and its lift to a triangulation $\tilde{\sigma}$ of the universal cover of ∂M . Note that each hyperbolic structure h_i on M defines a hyperbolic structure on the universal cover $\tilde{M}$ of M , and thus a polyhedral embedding of $\partial\tilde{M}$ in H^3 .

Definition 8.2. *Let S be a finite subset of the set of vertices of $\tilde{\sigma}$, with cardinal at least 2. We say that S **collapses** if there exists a subsequence of (h_i) and a sequence of conformal maps $\rho_i : H^3 \mapsto B_0(1) \subset \mathbf{R}^3$ such that:*

- *the set $\rho_i(S)$ converges to a point x_0 in S^2 ;*
- *no vertex of $\tilde{\sigma}$, adjacent to a vertex in S , converges to x_0 .*

Here $B_0(1)$ is the radius 1 ball centered at 0 in $\mathbf{R}^3$. This notion is useful because it implies the existence of a non-elementary circuit in σ on which the sum of the dihedral angles goes to 2π .

Proposition 8.3. *If a finite subset S of vertices of σ (with cardinal at least 2) collapses, then, maybe after taking a subsequence of (h_i) , there is a non-elementary circuit in the 1-skeleton of σ on which the sum of the dihedral angles goes to 2π .*

Proof. The argument is the same as in section 4 (and that in [Sch98a, Sch01a]): the polyhedron dual to $\partial\tilde{M}$ in de Sitter space, for the limit hyperbolic structure, has a face tangent to the boundary at infinity at the point where the collapse occurs, and the metric induced on this face is the metric of a hemisphere. Therefore the length of its boundary is 2π , and this provides a non-elementary circuit in $\tilde{\sigma}$ with sum of the dihedral angles converging to 2π . $\square$

Note that the non-elementary circuit on which the sum of the dihedral angles goes to 2π is simply the sequence of edges between (a connected component of) S and its complement.

We now have to define the kind of metrics that appear on the boundary of an ideal hyperbolic manifold M . Here S is a compact orientable surface of genus $g \geq 2$, with N points $v_1, \dots, v_N$ removed ($N \geq 1$), along with a triangulation σ whose vertices are the v_i — for instance, S can be ∂M with the vertices removed. We call $a_1, a_2, \dots, a_e$ the edges of σ , and $\tau_1, \dots, \tau_t$ its triangles.

Definition 8.4. *A hyperbolic structure on (S, σ) is a family of gluings between the adjacent triangles of σ , each being endowed with the metric of the ideal triangle in H^2 . The set of hyperbolic structures on (S, σ) will be denoted by $\mathcal{M}$.*

Obviously a hyperbolic structure in this sense defines a hyperbolic metric on S , but it is in general not complete; indeed, there might be a “shift” at some of the edges, in the sense described below.

Definition 8.5. *Let g be a hyperbolic structure on (S, σ) , and let E be an edge of σ . Choose an orientation of E , and let T_+ and T_- be the triangles of σ standing on the “right” and on the “left” of E respectively. Let u_+ and u_- be the orthogonal projections on E of the vertices opposite to E in T_+ and T_- respectively. The **shift** $sh(E)$ of g at E is the oriented distance, along E , between u_- and u_+ .*

Note that $sh(E)$ does not depend on the orientation chosen for E . This definition is somehow related to the notion of “shear” defined above for an ideal simplex at an edge, but it is better to use a different name to avoid confusions.

Proposition 8.6. *Suppose that no finite set of vertices of σ collapses. Then, after taking a subsequence, the shifts of the edges of σ for the h_i converge.*

Proof. Choose a “fundamental domain” in $\tilde{\sigma}$, i.e. a connected subgraph $\sigma_0 \subset \tilde{\sigma}$ such that each edge in σ has a unique inverse image in σ_0 for the canonical projection $\tilde{\sigma} \mapsto \sigma$.

We consider again the polyhedral embeddings $\phi_i : \partial\tilde{M} \rightarrow H^3$ associated to the ideal hyperbolic structures h_i on M . Since σ_0 has a finite number of edges, an elementary compactness argument shows that there exists a sequence (ρ_n) of conformal maps from H^3 to the radius 1 ball in $\mathbf{R}^3$ such that the images by $\rho_i \circ \phi_i$ of the vertices of σ_0 converge in S^2 .

Since by hypothesis there is no collapsing subset of vertices, there exists a constant $C > 0$ such that the edges of the image in S^2 of σ_0 by the limit have length between $1/C$ and C . An elementary argument then shows that the distance on S^2 between the vertices of σ_0 and the orthogonal projections on the edges of the triangles of the opposite vertices remain bounded between $1/C'$ and C' , where $C' > 0$ is another constant.

This in turns indicates that (after taking a subsequence) the shifts of σ converge. $\square$

Proof of lemma 8.1. By hypothesis the dihedral angles of the (h_i) converge, moreover the metrics induced on the boundary converge by proposition 8.6. Thus the polyhedral embeddings of $\partial\tilde{M}$ in H^3 converge, and therefore the hyperbolic structures (h_i) also do so. $\square$

8.2 Dihedral angles with a given triangulation

We now have all the tools necessary to describe the dihedral angles of ideal manifolds with a given triangulation of the boundary. We consider a triangulation Σ of M .

Lemma 8.7. *Consider a triangulation Σ of the interior of M . Let $\mathcal{H}_{\text{sm}}^\Sigma$ be the set of smooth hyperbolic structure obtained by gluing ideal simplices according to Σ . Then:*

1. *the space of possible dihedral angles forms the interior of a convex polyhedron Θ_Σ in $\mathbf{R}^{e_b}$.*
2. *for any $\theta \in \Theta_\Sigma$, there is a unique smooth hyperbolic structure on M with those dihedral angles; $\mathcal{H}_{\text{sm}}^\Sigma$ therefore inherits the affine structure of Θ_Σ .*
3. *on each point of the boundary of Θ_Σ , one of the following happens:*
 - (a) *one of the simplices of Σ has a dihedral angle which goes to 0;*
 - (b) *the dihedral angle of one of the boundary edges goes to 0 or to π ;*
 - (c) *there is a non-elementary circuit in the 1-skeleton of Σ for which the sum of the dihedral angles goes to 2π .*
4. *the volume V is a smooth, strictly concave function on Θ_Σ .*

The proofs of points (1) and (2) are consequences of the previous section. Indeed the condition that the dihedral angles are given by θ defines a family of linear conditions on the angles, so an affine subspace of the space of angle assignations of the simplices of Σ . Adding the condition that the excess angle at each interior edge is 2π defines an affine subspace, and thus a smaller polyhedron in the space of angle assignations on the simplices of Σ . Projecting to the space of possible dihedral angles on the boundary edges thus defines a convex polyhedron.

Point (2) follows from a deformation argument. Consider the map $\Phi : \mathcal{H}_{\text{sm}}^\Sigma \rightarrow \Theta_\Sigma$ sending an ideal hyperbolic structure to its dihedral angles. Corollary 7.11 shows that Φ is locally injective (its differential is injective), while lemma 8.1 shows that Φ is proper — it is therefore a covering. But $\mathcal{H}_{\text{sm}}^\Sigma$ is connected, while Θ_Σ is contractible, so Φ is a homeomorphism. Therefore each assignation of dihedral angles in Θ_Σ is indeed realized by a smooth hyperbolic structure.

Point (3) is a consequence of lemma 8.1, which indicates that no degeneration other than (a), (b) and (c) can occur.

Point (4) is also a consequence of the definitions, and of the following remarks, which applies directly to this case.

Remark 8.8. *Let $\Omega \in \mathbf{R}^N$ be a convex subset, and let $f : \Omega \rightarrow \mathbf{R}$ be a smooth, strictly concave function. Let $\rho : \mathbf{R}^N \rightarrow \mathbf{R}^p$ be a linear map, with $p < N$, and let $\overline{\Omega} := \rho(\Omega)$. Define a function:*

$$\begin{aligned} \overline{f} : \overline{\Omega} &\rightarrow \mathbf{R} \\ y &\mapsto \max_{x \in \rho^{-1}(y)} f(x) \end{aligned}$$

Then $\overline{\Omega}$ is convex, and $\overline{f}$ is a smooth, strictly concave function on $\overline{\Omega}$.

Proof. It is quite obvious that $\overline{\Omega}$ is convex, and also that $\overline{f}$ is smooth since f is strictly concave.

Let $\overline{c} : [0, 1] \rightarrow \overline{\Omega}$ be a geodesic segment, parametrized at constant speed. By definition of $\overline{f}$, there exist points $x_0, x_1 \in \Omega$ such that:

$$\overline{c}(0) = \rho(x_0), \quad \overline{c}(1) = \rho(x_1), \quad \overline{f} \circ \overline{c}(0) = f(x_0), \quad \overline{f} \circ \overline{c}(1) = f(x_1) .$$

Let $c : [0, 1] \rightarrow \Omega$ be the geodesic segment parametrized at constant speed such that $c(0) = x_0$ and $c(1) = x_1$. Since ρ is linear, $\overline{c} = \rho \circ c$.

Moreover, since f is strictly concave

$$\forall t \in (0, 1), \quad f \circ c(t) > tf \circ c(0) + (1 - t)f \circ c(1) .$$

Therefore, the definition of $\overline{f}$ shows that:

$$\forall t \in (0, 1), \quad \overline{f} \circ \overline{c}(t) \geq f \circ c(t) > tf \circ c(0) + (1 - t)f \circ c(1) = \overline{f} \circ \overline{c}(0) + \overline{f} \circ \overline{c}(1) .$$

This shows that $\overline{f}$ is strictly concave. □

8.3 Fixed triangulations of the boundary

Using the ideas and results of the previous section, we can now give a description of the set of dihedral angles which can be achieved on the ideal hyperbolic manifolds with a given cellulation of the boundary. Studying the possible degenerations and a global argument using the results of section 3 (on the fuchsian case) will then lead to theorem 8.15. So we now consider an ideal hyperbolic manifold M , which does not necessarily admit an ideal triangulation. Of course the point is that, by lemma 5.4, a finite cover of M admits one.

Lemma 8.9. *Consider a cellulation σ of the boundary ∂M , and let $\mathcal{H}_{sm}^\sigma$ be the set of ideal hyperbolic structures on M such that the boundary is triangulated according to σ . Then each connected component of $\mathcal{H}_{sm}^\sigma$ has a natural affine structure $\mathcal{A}_\sigma$, obtained by gluing the $\mathcal{H}_{sm}^\Sigma$ — for Σ a triangulation of the interior of M inducing the triangulation σ of the boundary — in the natural way on their intersections. $(\mathcal{H}_{sm}^\sigma, \mathcal{A}_\sigma)$ is affinely equivalent to the disjoint union of the interiors of a finite set of convex polyhedra in $\mathbf{R}^{eb}$, and V is a concave function on each.*

Proof. Let $\theta \in \Theta_\sigma$, and let $h \in \mathcal{H}_{sm}^\sigma(\theta)$. By lemma 7.14, the infinitesimal deformations of h are parametrized by the admissible infinitesimal deformations of the dihedral angles at the exterior edges.

Let Θ_Σ be the set of dihedral angles that can be attained by deformation from h . By proposition 8.10, the boundary is made of angle assignments satisfying one of a number of possible affine equalities, so that Θ_Σ is the interior of a convex polyhedron in $\mathbf{R}^{eb}$. Moreover, its boundary can be decomposed into the union of two components, $\partial\Theta_\Sigma = \partial_t\Theta_\Sigma \cup \partial_r\Theta_\Sigma$, where:

1. $\partial_r\Theta_\Sigma$ is the set of boundary points where the sum of the dihedral angles on the edges of a non-elementary circuit goes to 2π , or the dihedral angle at some boundary edge goes to 0 or to π .
2. $\partial_t\Theta_\Sigma$ is the set of boundary points where the previous condition does not apply, but one of the simplices has an angle which goes to 0.

By lemma 3.11, $\partial_r\Theta_\Sigma$ corresponds to the boundary of $\mathcal{H}_{sm}^\sigma$. On the other hand, let $\bar{h}$ correspond to a point of $\partial_t\Theta_\Sigma$. Lemma 7.14 shows that $\bar{h}$ can be deformed so that its dihedral angles go beyond $\partial_t\Theta_\Sigma$. So there is another ideal triangulation Σ' of $(M, \bar{h})$ such that $\bar{h} \in \mathcal{H}_{sm}^{\Sigma'}$. Since the affine structure on $\mathcal{H}_{sm}^\Sigma$ is defined in terms of the variations of the dihedral angles, it extends naturally to an affine structure on $\mathcal{H}_{sm}^{\Sigma'}$. This shows that $\mathcal{H}_{sm}^\sigma$ carries an affine structure $\mathcal{A}_\sigma$, obtained by gluing the affine structures on the $\mathcal{H}_{sm}^\Sigma$.

Moreover, at a point of $\partial_t\Theta_\Sigma \cap \partial_r\Theta_\Sigma$, the faces of $\mathcal{H}_{sm}^{\Sigma'}$ corresponding to $\partial_r\Theta_\Sigma$ are the extensions of the faces of $\mathcal{H}_{sm}^\Sigma$, so that $\mathcal{H}_{sm}^\sigma$ is locally convex for the affine structure. It is therefore affinely equivalent to the interior of a convex polyhedron in $\mathbf{R}^{eb}$. $\square$

In addition, lemma 8.7 allows for a simple description of the possible boundary behavior; since the cases corresponding to the collapse of a simplex correspond to the boundary of the cells $\mathcal{H}_{sm}^\Sigma$ rather than to the boundary of $\mathcal{H}_{sm}^\sigma$, the boundary of $\mathcal{H}_{sm}^\sigma$ is characterized by one of the other possible limit cases.

Proposition 8.10. *Let $(h_n)_{n \in \mathbf{N}}$ be a sequence of elements of $\mathcal{H}_{sm}^\sigma$ converging to a limit $h \in \partial\mathcal{H}_{sm}^\sigma$. After taking a subsequence, one of the following occurs:*

1. *some boundary edge has a dihedral angle which goes to 0 or π ;*
2. *there exists a non-elementary circuit in $\Sigma_{|\partial M}$ on which the sum of the dihedral angles goes to 2π .*

8.4 Spaces of angle assignments

The results already obtained give us a good understanding of the local properties of the map sending an ideal hyperbolic manifold to the angle assignment on the boundary edges. To complete the proof of theorem 8.15, we need some global topological information on the space of angle assignments on the boundary satisfying the conditions of the theorem.

We need some additional notations. We will call $(\partial_i M)_{1 \leq i \leq N}$ the connected components of ∂M . For each $n \in (\mathbf{N} \setminus \{0\})^N$, we call $\mathcal{A}_n$ the space of couples (Γ, w) , where Γ is the 1-skeleton of a cellulation of

∂M having n_i vertices in each $\partial_i M$, and $w : \Gamma_1 \rightarrow (0, \pi)$ is a function satisfying the conditions of theorem 8.15; and we call $\mathcal{M}_n$ the space of ideal hyperbolic manifolds having n_i vertices in $\partial_i M$, for each i . Φ_n will then be the natural map sending an element of $\mathcal{M}_n$ to the corresponding dihedral angles assignation on a cellulation of ∂M , which is in $\mathcal{A}_n$.

We will prove that if M has incompressible boundary, then, for each n , $\mathcal{A}_n$ is connected. Thus Φ_n is a covering. To prove that it is actually a homeomorphism, we will consider some special dihedral angle assignations for which the uniqueness follows from the Mostow rigidity.

Lemma 8.11. *Suppose that M has incompressible boundary. Then, for each $n \in (\mathbb{N} \setminus \{0\})^N$, $\mathcal{A}_n$ is connected.*

This is the only point where we use the fact that M has incompressible boundary.

Recall that, for $g \geq 2$ and $r \geq 1$, $\mathcal{A}_{g,r}$ is the space of dihedral angles assignations for fuchsian manifolds of genus g , with r vertices. From now on, we call g_i the genus of $\partial_i M$. With the topological assumptions which we have made on M , $g_i \geq 2$ for each i .

Remark 8.12. *Suppose that M has incompressible boundary. Then, for each $n \in (\mathbb{N} \setminus \{0\})^N$, $\mathcal{A}_n$ is homeomorphic to $\prod_{i=1}^N \mathcal{A}_{g_i, n_i}$.*

Proof. The conditions in theorem 8.15, which define both $\mathcal{A}_n$ and the $\mathcal{A}_{g_i, n_i}$, describe each connected component of ∂M independently from the others. Since M has incompressible boundary, the curves in each connected component of ∂M which are contractible in M are those which are contractible in ∂M ; so the condition on the dihedral angles on $\partial_i M$ are exactly those describing $\mathcal{A}_{g_i, n_i}$. $\square$

Proposition 8.13. *For each $g \geq 2$ and each $r \geq 1$, $\mathcal{A}_{g,r}$ is homeomorphic to the space $\mathcal{T}_{g,r}$ of conformal structures on a surface of genus g , with r marked points (up to isotopy).*

Proof. Let Σ_g be a compact surface of genus g , and let t be a conformal structure on Σ_g . There is a unique hyperbolic metric on Σ_g , say h , in the conformal class defined by t . Taking the warped product:

$$M_g := (\Sigma_g \times \mathbb{R}, dt^2 + \cosh^2(t)h)$$

determines a complete fuchsian hyperbolic manifold. If $p_1, \dots, p_r$ are points in Σ_g , they define points $p_{1,\pm}, \dots, p_{r,\pm}$ in $\partial_\infty M_g$, with $p_{i,+}$ in the “upper” boundary and $p_{i,-}$ in the “lower” boundary for each i , and $p_{i,+}$ exchanged with $p_{i,-}$ by the isometric involution on M_g .

Let $\overline{M}$ be the convex hull of the $p_{i,\pm}$. By remark 3.9, $\overline{M}$ is an ideal hyperbolic manifold. Moreover, each fuchsian ideal hyperbolic manifold is obtained in this way, and thus is associated to a $t \in \mathcal{T}_{g,r}$.

Finally, theorem 4.25 shows that $\mathcal{A}_{g,r}$ is homeomorphic to the space of fuchsian ideal hyperbolic manifolds with each boundary surface of genus g with r vertices, and the result follows. $\square$

Proof of lemma 8.11. For each $g \geq 2$ and $r \geq 1$, $\mathcal{T}_{g,r}$ is connected. So the result follows from the previous proposition. $\square$

8.5 Unique realization for some dihedral angle assignations

The content of the previous subsection is sufficient to ensure that the map Φ_n is a covering for each choice of n , and that the number of inverse images of all elements of $\mathcal{A}$ is the same. The next lemma states that this number is 1 for some cellulations of M and choices of dihedral angles of the edges.

Lemma 8.14. *Let σ be a cellulation of ∂M such that:*

- *the faces of σ can be separated into two sets F_- and F_+ , with each edge bounding a face of F_- and one of F_+ .*
- *each vertex is adjacent to 4 faces.*

Let $w : \sigma_1 \rightarrow (0, \pi)$ be the function assigning the value $\pi/2$ to each edge. Then w satisfies the hypothesis of theorem 8.15, and there is at most one ideal hyperbolic manifolds with combinatorics given by σ and dihedral angles given by w .

Proof. Let g be a ideal hyperbolic metric on M with boundary combinatorics σ and all angles $\pi/2$. We consider 4 copies of M , say M_1, M_2, M_3 and M_4 , each with the metric g . We identify the corresponding faces of the M_i according to the following table, where a “+” on line i and column j means that each face of M_i in F_+ is glued to the corresponding face on M_j , and the same for $-$.

	1	2	3	4
1		-		+
2	-		+	
3		+		-
4	+		-	

Let N be the result of those gluings. It is easy to check that:

1. the topology of N is determined by the topology of M and of σ , and does not depend on the ideal hyperbolic metric on M .
2. N , with the metric coming from the gluing of the M_i , is a complete, finite volume hyperbolic manifold.
3. two non-isometric ideal hyperbolic metrics on M would result in two non-isometric hyperbolic metrics on N .

By Mostow rigidity there is at most one complete, finite volume hyperbolic metric on N . The result follows. $\square$

8.6 Proof of the main theorem

We can now prove theorem 8.15, concerning the description of dihedral angles of ideal hyperbolic manifolds.

Theorem 8.15. *Suppose that M has incompressible boundary. Then the set Θ_σ of dihedral angles of ideal hyperbolic structures whose boundary is cellulated according to σ is given by the condition that:*

1. on each elementary circuit in σ , the sum of the exterior dihedral angles is 2π ;
2. on each non-elementary circuit, the sum of the dihedral angles is strictly larger than 2π .

Each dihedral angles assignation is obtained on a unique ideal hyperbolic manifold.

Proof. Let σ be a cellulation of ∂M . By lemma 8.9, there exists a number $\nu(\sigma) \in \mathbf{N}$ such that each dihedral assignation on the edges of σ , satisfying the hypothesis of the theorem, is realized by exactly $\nu(\sigma)$ ideal hyperbolic manifolds with boundary combinatorics σ .

Let $n \in (\mathbf{N} \setminus \{0\})^N$. By lemma 8.11, $\mathcal{A}_n$ is connected; therefore, for all cellulations σ of ∂M with n_i vertices in $\partial_i M$, the value of $\nu(\sigma)$ is the same, and is equal to a number $\nu(n) \in \mathbf{N}$.

Let σ be a cellulation of ∂M . Let $\bar{\sigma}$ be the cellulation of ∂M obtained by “splitting” a vertex v of σ , i.e. replacing v by two vertices v_1 and v_2 connected by an edge e , and replacing two edges e_1 and e_2 containing v by two edges each, say e_{11}, e_{12} and e_{21}, e_{22} . If $v \in \partial_i M$, let $\bar{n} = (n_1, n_2, \dots, n_{i+1}, \dots, n_N)$.

Each dihedral angle assignation θ to the edges of σ can be identified with a “degenerate” dihedral angle assignation $\bar{\theta}$ on $\bar{\sigma}$, with a dihedral angle 0 at e , dihedral angles at e_{i1} and e_{i2} equal to half the dihedral angles at e_i , and the other dihedral angles equal to those of θ . If θ satisfies the condition of theorem 8.15, so does $\bar{\theta}$ (except of course for the fact that one of its angles is 0). Let (θ_n) be a sequence of dihedral angle assignations on the edges of $\bar{\sigma}$ which converges to $\bar{\theta}$.

For each $i \in \{1, \dots, \nu(\bar{n})\}$, there is a sequence of ideal hyperbolic metrics $(g_{i,n})_{n \in \mathbf{N}}$ with dihedral angles equal to θ_n . As $n \rightarrow \infty$, $(g_{i,n})$ converges to an ideal hyperbolic metric g_i on ∂M with boundary combinatorics given by σ , and dihedral angles by θ . So $\nu(\bar{n}) \geq \nu(n) \geq 1$.

But it is easy to check that, for each $n \in (\mathbf{N} \setminus \{0\})^N$, there exists some $p \in (\mathbf{N} \setminus \{0\})^N$, with $p_i \geq n_i$ for all $i \in \{1, \dots, N\}$, such that there exists a cellulation of ∂M , with p_i vertices in each $\partial_i M$, to which lemma 8.14 applies.

So, for each $n \in (\mathbf{N} \setminus \{0\})^N$, $\nu(n) = 1$, which proves the theorem. $\square$

It might be useful to note that the convexity of the boundary of M is not really necessary for many steps of the proof of this theorem; for instance the infinitesimal rigidity results should still hold for at least some manifolds with non-convex boundary. This contrasts with most results concerning the induced metric on the boundary, where the convexity is crucial.

8.7 An affine structure on some Teichmüller spaces

A consequence of the previous considerations is that there is a natural affine structure on the space of ideal hyperbolic structures on M , coming from the parametrization by the dihedral angles. Its definition is simple and depends directly only on the dihedral angles. It has “cone-like” singularities along the “cells” of codimension at least 2, corresponding to non-generic polyhedral structures of the boundary. The remarkable point, however, is that it also has a well-defined volume element. This, which I can not explain better than by doing an explicit computation, might indicate that this affine structure is somehow meaningful. The definition below should be easier to understand with figure 4.

Definition 8.16. *The affine structure $\mathcal{A}$ on $\mathcal{H}_{sm}$ is defined as follows.*

- for each triangulation σ of the boundary, the restriction of $\mathcal{A}$ to $\mathcal{H}_{sm}^\sigma$ is defined as $\mathcal{A}_\sigma$ above.
- if σ is a cellulation with one 4-gonal face f and the other faces triangles, then $\mathcal{H}_{sm}^\sigma$ is a codimension 1 face of $\mathcal{H}_{sm}$ which bounds two maximal dimension faces $\mathcal{H}_{sm}^{\sigma_1}$ and $\mathcal{H}_{sm}^{\sigma_2}$, where σ_1, σ_2 are triangulations obtained by refining σ by adding edges e_1 and e_2 respectively to the 4-gonal face of σ . Then $\mathcal{H}_{sm}^{\sigma_1}$ and $\mathcal{H}_{sm}^{\sigma_2}$ are glued by the map, from a neighborhood of σ in the extension of σ_1 corresponding to having a negative angle at e_1 to $\mathcal{H}_{sm}^{\sigma_2}$, which sends a configuration with angle $-2u$ at e_1 and a, b, c, d at the 4 edges of f , to a configuration with angle $2u$ at e_2 , and angles $a-u, b-u, c-u, d-u$ at the edges of f .




Figure 4: Definition of $\mathcal{A}$, on the dual graph

Note that this definition makes sense since the transformation preserves the angle conditions defining the space of possible angle assignments. It is clear that the lengths conditions on elementary and non-elementary circuits are satisfied for one of the configurations if and only if they are satisfied for the other.

Lemma 8.17. *$\mathcal{A}$ has a holonomy preserving a volume form.*

Proof. To prove this one should check that the holonomy around all codimension two faces of $\mathcal{H}_{sm}$ preserve a volume form. There are two kinds of codimension two faces, those corresponding to a cellulation of ∂M with two 4-gonal faces, and those corresponding to the cellulations with one 5-gonal face. In the first case of the holonomy is trivial, so the only case to check is the second.

When a cellulation σ of ∂M has one 5-gonal face and its other faces are triangles, the corresponding cell $\mathcal{H}_{sm}^\sigma$ of $\mathcal{H}_{sm}$ bounds five maximal dimension and five codimension one faces. The holonomy around $\mathcal{H}_{sm}^\sigma$ is computed by considering the diagram in figure 5, corresponding to the transformation of the angles as one goes through the five maximal dimension faces — the angles are marked on the dual graph, and the constant part of the angles on the five “exterior” edges are not marked.




Figure 5: Holonomy computation

The only part of the holonomy which is relevant for the volume is the one corresponding to the two “central” edges. The computation made on figure 5 shows that the holonomy for those two angles is given by the matrix:

$$M := \frac{1}{32} \begin{pmatrix} 28 & -10 \\ 10 & 33 \end{pmatrix}$$

But $\det(M) = 1$, so that the holonomy around $\mathcal{H}_{\text{sm}}^\sigma$ preserves a volume form. $\square$

Remark 8.18. $(\mathcal{H}_{\text{sm}}, \mathcal{A})$ has locally convex boundary at all points which are in the closure of a face $\mathcal{H}_{\text{sm}}^\sigma$.

Proof. We have to prove that if σ is a triangulation of ∂M and F_0 is a codimension 1 face of $\mathcal{H}_{\text{sm}}^\sigma$ which is in the boundary of $\mathcal{H}_{\text{sm}}$, then, across each codimension 1 face of $\mathcal{H}_{\text{sm}}^\sigma$ which is not in $\partial \mathcal{H}_{\text{sm}}$, the extension of F_0 is either outside $\mathcal{H}_{\text{sm}}$ or in its boundary.

Let $\mathcal{H}_{\text{sm}}^{\sigma_1}$ be a codimension 1 face of $\mathcal{H}_{\text{sm}}^\sigma$, so that σ_1 is obtained from σ by removing an edge from σ , and σ_1 has exactly one face f with 4 edges. There are 3 cases to consider:

1. F_0 corresponds to a non-elementary circuit on which the sum of the dihedral angles is 2π . Then the definition of $\mathcal{A}$ shows directly that the extension of F_0 beyond $\mathcal{H}_{\text{sm}}^{\sigma_1}$ remains in the boundary of $\mathcal{H}_{\text{sm}}$.
2. F_0 corresponds to an edge $e \notin f$ with dihedral angle equal to π . It is then clear again from the definition that the extension of F_0 beyond $\mathcal{H}_{\text{sm}}^{\sigma_1}$ remains in $\partial \mathcal{H}_{\text{sm}}$, since the transformation which occurs in the definition of $\mathcal{A}$ does not change the angle at e .
3. F_0 corresponds to an edge $e \in f$ with dihedral angle equal to π . But then the definition of $\mathcal{A}$ shows that, if one goes beyond F_0 on an affine line starting in $\mathcal{H}_{\text{sm}}^{\sigma_1}$, the angle at e is above π after $\mathcal{H}_{\text{sm}}^{\sigma_1}$, so that the extension of F_0 beyond $\mathcal{H}_{\text{sm}}^{\sigma_1}$ is outside $\mathcal{H}_{\text{sm}}$.

$\square$

An elementary consequence of the constructions which we have just seen — applied to the fuchsian case — is that there is a natural affine structure on the Teichmüller space of a surface of genus at least 2, with at least one marked point. The point is that fuchsian manifolds form an affine submanifold of the ideal hyperbolic manifolds (they corresponds to the case where the triangulations and dihedral angles are the same on the two components of the boundary). Moreover, if one remains in the category of fuchsian manifolds, there is no bent case (see remark 3.9), so that the affine structure defined on ideal manifolds spans the whole Teichmüller space. One thus obtain the:

Theorem 8.19. *For each $g \geq 2$ and each $N \geq 1$, there is a natural unimodular piecewise affine structure $\mathcal{A}_{g,N}$ on the Teichmüller $\mathcal{T}_{g,N}$ space of the genus g surface with N marked points.*

There are some natural questions that remain on this structure; for instance, how many cells it has, and whether the affine structure is globally convex.

8.8 Hyperbolic polyhedra

The previous results apply in a specially simple way to ideal polyhedra in H^3 , i.e. the case where M is topologically a ball. Then, for a fixed number $N \geq 3$ of vertices, there is a fixed number of combinatorial types of possible Delaunay cellulations of S^2 with N vertices, so that the space $\mathcal{H}_{\text{sm}}$ of dihedral angles assignments is the union of a finite number of cells. But even here some questions remains, for instance whether the set of ideal polyhedra with N vertices, with the affine structure coming from the dihedral angles, is globally convex.

9 Induced metrics

We will study in this section some properties of the metrics induced on the boundaries of ideal hyperbolic manifolds. They are finite area hyperbolic metrics on each connected component of the boundary of M without its ideal points, and we will show that the infinitesimal deformations of the interior metric are parametrized by the infinitesimal deformations of the boundary metric; this is again a consequence of the Schläfli formula. In addition, we will prove a global result in the special case of fuchsian manifolds, then the induced metrics on the boundary are in one-to-one correspondence with the hyperbolic structures on the interior.

9.1 Finite area metrics on surfaces

We will use again here the notion of hyperbolic structure on a triangulated surface, as defined in the previous section. We consider again a compact, orientable surface S of genus at least 2 with N points $v_1, \dots, v_N$ removed. The following additional definition is natural.

Definition 9.1. *Let $g \in \mathcal{M}$ and let v be a vertex of σ . The **shift** of g at v is the sum of the shifts of g at the edges containing v . We will say that g is **complete** if its shift is zero at all vertices. The set of complete structures will be denoted by $\mathcal{M}_c$.*

Of course the notion of completeness defined here is the same as the usual, topological notion. Indeed if the shift of g at a vertex v is non-zero, it is possible to use this — and the fact that ideal triangles are exponentially thin near their vertices — to attain v in a finite time, by “circling” around it to take opportunity of the shift. The reciprocal is not difficult to prove either.

Finally, knowing the shifts of a complete metric $g \in \mathcal{M}_c$ at all edges determines all the gluing along the edges; let S_0 be the set of maps from the set of edges of σ to $\mathbf{R}$ such that, at each vertex, the sum of the adjacent edges is 0. We identify S_0 with $\mathbf{R}^{e-v}$. Then:

Proposition 9.2. *The map F from $\mathcal{M}_c$ to $S_0 = \mathbf{R}^{e-v}$ sending g to $F(g)$ defined, for an edge E , by $F(g)(E) = sh_g(E)$, is a bijection.*

Proof. It is quite obvious that the hyperbolic structures on a given triangulated surface are determined by their shifts. On the other hand each shift function on the edges corresponds to a possible gluing (under the condition that the sum on the edges at any vertex is zero, by completeness). $\square$

9.2 The lengths of the edges of σ

Let $g \in \mathcal{M}_c$. Each vertex v_i of σ has a neighborhood which is isometric to a neighborhood of a cusp in the quotient of H^2 by a parabolic isometry. For each such v_i , choose a horocycle H_i centered at v_i . Then, for each edge E of σ , going from a vertex v_i to a vertex v_j , let $l(E)$ be the oriented length of E between H_i and H_j . The orientation is chosen so that $l(E)$ is negative when the horoballs bounded by H_i and H_j overlap. Clearly, replacing H_i by another horocycle centered at v_i changes the lengths of the edges containing v_i as an end by the addition of a constant. So l defines a function:

$$l : \mathcal{M}_c \rightarrow \mathbf{R}^e / \mathbf{R}^v .$$

Proposition 9.3. *l is a bijection between $\mathcal{M}_c$ and $\mathbf{R}^e / \mathbf{R}^v$.*

The proof uses the following elementary property of ideal triangles in H^2 .

Sub-lemma 9.4. *Let x_1, x_2, x_3, x_4 be four distinct points on $S^1 = \partial_\infty H^2$, in this cyclic order. For each $i \in \{1, 2, 3, 4\}$, let h_i be a horocycle centered at v_i , and, for $i \neq j$, let l_{ij} be the distance between h_i and h_j along the geodesic going from v_i to v_j — which is negative if h_i and h_j overlap. Let π_2 and π_4 be the orthogonal projections on (x_3, x_1) of x_2 and x_4 respectively, and let δ be the oriented distance between π_2 and π_4 on (x_3, x_1) . Then:*

$$2\delta = l_{12} - l_{23} + l_{34} - l_{41} .$$




Figure 6: Ideal triangles (in the projective model of H^2)

Proof. It follows from figure 6, where numbers from 1 to 6 are attached to lengths of segments. Elementary properties of the ideal triangle show that:

$$\begin{aligned} l_{12} - l_{23} + l_{34} - l_{41} &= (1 + 2) - (2 + 3) + (4 + 5) - (5 + 6) \\ &= 1 - 3 + 4 - 6 \\ &= (1 - 6) + (4 - 3) \\ &= 2\delta . \end{aligned}$$

□

Proof of proposition 9.3. Using proposition 9.2, we are brought to proving that the map:

$$L : S_0 \simeq \mathbf{R}^{e-v} \rightarrow \mathbf{R}^e / \mathbf{R}^v$$

sending the e -uple of shifts of a metric to the lengths of its edges is a bijection. But the sub-lemma above shows that L is linear, and explicitly describes its inverse. □

9.3 Rigidity and the Schläfli formula

We now consider an ideal hyperbolic manifold M ; thanks to the previous paragraphs, we know that, if ∂M is triangulated, then its induced metric is determined by the lengths of the edges (defined up to the addition of a constant for each vertex). Of course, if ∂M is not triangulated — i.e. if some of the faces are polygons with more than three vertices — one can subdivide its cellulation to obtain a triangulation.

We will prove in this paragraph and the next that M is infinitesimally rigid, in the following sense. We call Φ the map from $\mathcal{H}_{\text{sm}}$ to $\mathcal{M}_c$ sending $g_0 \in \mathcal{H}_{\text{sm}}$ to the metric induced on its boundary. Then:

Lemma 9.5. *For any $g_0 \in \mathcal{H}_{\text{sm}}$, $T_{g_0}\Phi$ is an isomorphism.*

The proof will take two steps. We first treat the case when the boundary of M is triangulated, then the general case. The first case will rely on the Schläfli formula and remark 8.8. Recall from lemma 8.7 that, for any ideal hyperbolic manifold (M, g_0) , the volume V is a strictly concave function on $\mathcal{H}_{\text{sm}}$ in the neighborhood of g_0 .

Proposition 9.6. *Suppose that all faces of (M, g_0) are triangles, and that it is triangulable. Then $T_{g_0}\Phi$ is an isomorphism.*

Proof. We know by corollary 7.13 that $\dim \mathcal{H}_{\text{sm}} = e - v = \dim \mathcal{M}_c$. Moreover, the volume $V(g)$ is a strictly concave function on $\mathcal{H}_{\text{sm}}$ by lemma 8.7.

Recall the Schläfli formula from section 1:

$$dV = -\frac{1}{2} \sum_E l(E) d\theta(E) ,$$

where the sum is over all edges, and $\theta(E)$ is the interior dihedral angle of edge E . The strict concavity of V with respect to the $\theta(E)$ means that the Hessian of V with respects to those angles is negative definite, so that the differential of the lengths $l(E)$ — with respect to the dihedral angles — is non-degenerate. Thus $T_g l$ is injective, and the result follows. $\square$

Now let (M, g) be an ideal hyperbolic manifold with non-triangular boundary. By lemma 5.4, M has a finite covering $\overline{M}$ which admits an ideal triangulation Σ . Σ induces an ideal triangulation of $\partial \overline{M}$, and the argument above — the convexity of the volume and the Schläfli formula — still implies that, for each assignation of a real number $r(e)$ to each edge e of $\partial \overline{M}$ (defined up to addition of a constant for each vertex) there is a unique infinitesimal deformation of $\overline{M}$ such that the length of e (defined with respect to a set of horospheres at the vertices) varies at the rate $r(e)$. Therefore the argument given above shows that the infinitesimal deformations of $\overline{M}$ are uniquely determined by the induced variation of the boundary metric, and that each variation of the boundary metric is obtained.

This is in particular true for the variation of the boundary metric which are invariant under the action of $\pi_1 M / \pi_1 \overline{M}$ on $\overline{M}$, and the uniqueness implies that they are associated to infinitesimal deformations of $\overline{M}$ equivariant under the same action. So we obtain the proof of lemma 9.5, which we can reformulate as:

Lemma 9.7. *For each ideal hyperbolic manifold M , and for each infinitesimal deformation $\overline{h}$ of the boundary metric h on ∂M , there is a unique infinitesimal deformation of M inducing $\overline{h}$.*

9.4 The fuchsian case

Lemma 9.7 provides only an infinitesimal deformation result for the induced metrics on the boundaries of ideal hyperbolic manifolds. It is then difficult to obtain a global existence and uniqueness results for the boundary case, in particular because we have no such infinitesimal rigidity result for bent hyperbolic manifolds, which necessarily enter the picture. But in the fuchsian case, the bent case is excluded by remark 3.9, and a global result can be achieved.

Theorem 9.8. *Let S be a surface of genus $g \geq 2$, and let $N \geq 1$. For each complete, finite area hyperbolic metric h on S with N cusps, there is a unique ideal fuchsian hyperbolic manifold M such that the induced metric on each component of the boundary is h .*

Proof. We will use a deformation proof, following the original approach of Aleksandrov [Ale58] for similar polyhedral question. We choose $g \geq 2$ and $N \geq 1$. We call $\mathcal{T}_{g,N}$ the Teichmüller space of marked conformal structures on the compact surface Σ_g of genus g with N marked points. There is a natural map $\Phi_{g,N}$ from $\mathcal{T}_{g,N}$ to itself, defined as follows. For $h \in \mathcal{T}_{g,N}$, the conformal class on Σ_g (i.e. forgetting the marked points on Σ_g) contains a unique hyperbolic metric h_0 on Σ_g ; taking its universal cover defines a conformal map from $\tilde{\Sigma}_g$ with the conformal structure in h to the upper hemisphere S^2_+ . The marked points in h then define an equivariant set S of points in S^2_+ . Taking the boundary of the convex hull of those points in $\mathbf{R}^3$ (and subtracting its intersection with the plane containing the equator) leads to a polyhedral surface, which is invariant under the natural action of $\pi_1 \Sigma_g$ on H^3 . The quotient is homeomorphic to Σ_g , and has N vertices (which are the intersection points with S^2_+); it carries a complete, finite area hyperbolic metric $\bar{h}$ induced by the canonical metric on H^3 . $\Phi_{g,N}(h)$ is defined as this metric $\bar{h}$, considered as an element of $\mathcal{T}_{g,N}$.

The proof of the theorem is an immediate consequence of the following points.

1. $\Phi_{g,N}$ is locally injective, i.e. its differential is an isomorphism at each point.
2. $\Phi_{g,N}$ is proper.
3. $\mathcal{T}_{g,N}$ is connected and simply connected.

Point (1) is just lemma 9.5, while point (3) is well known (since we use the marked Teichmüller space). So we only have to prove point (2). It can be reformulated as a degeneration statement: if $(M_n)_{n \in \mathbf{N}}$ is a sequence of fuchsian ideal manifolds of genus n with N vertices, and if the sequence of induced metrics (h_n) converges to a limit $h \in \mathcal{T}_{g,N}$, then (M_n) converges.

In other terms, we have to prove that if either the conformal structure on $\partial E(M_n)$ degenerates or the marked points collapse, then the length of some closed geodesic on ∂M goes to 0.

In the case of a degeneration of the conformal structure on $\partial E(M_n)$, the action of $\pi_1 \partial M$ on S^2_+ also degenerates, and it is then easy to check that some closed geodesic goes to 0. Similarly, if the set of marked points collapses, then the collapsing subset is bounded, in ∂M , by a long thin tube of very small diameter, and the compactness result follows. $\square$

9.5 Ideal polyhedra in H^3

An elementary remark is that, for ideal polyhedra in H^3 , the approach above gives a simple proof of the existence and uniqueness of ideal polyhedra having a given induced metric, see [Riv92, Riv96]. In particular this approach does not use the Cauchy method to get the global uniqueness, since it follows from the global deformation result.

Rather, the infinitesimal rigidity can be obtained as in lemma 9.7 using the Schläfli formula and the results concerning the dihedral angles. Applying a deformation argument to conclude then only requires a compactness result — namely, that if a sequence of ideal polyhedra has induced metrics which converge, then it has a converging subsequence (modulo isometries). Such a compactness result is easy to obtain directly.

10 Circles packings

10.1 From dihedral angles to circle configurations

As mentioned in section 2, the polyhedral questions considered here can be formulated in terms of configurations of circles in S^2 . More precisely, let M be an ideal hyperbolic manifold. Consider its universal cover $\tilde{M}$ as a subset of H^3 . Its boundary $\partial \tilde{M}$ is the disjoint union of a set of convex surfaces, and their boundary is the limit set Λ of the action of $\pi_1 M$ on H^3 .

Each face of $\tilde{M}$ defines an oriented totally geodesic plane in H^3 , with boundary at infinity an oriented circle in $S^2 \setminus \Lambda$. If two faces F, F' of $\partial \tilde{M}$ are adjacent, then the corresponding oriented circles C, C' intersect with angle equal to the exterior dihedral angle between F and F' . Moreover, a simple convexity argument shows that the union of the interiors of the circles corresponding to the faces of $\partial \tilde{M}$, along

with the intersection points, is $S^2 \setminus \Lambda$. So the results concerning the dihedral angles of ideal hyperbolic manifolds can be formulated in terms of configurations of circles in the complement of Λ , with given intersection angles.

Therefore, an ideal hyperbolic manifold determines a circle configuration on ∂M , in the following sense.

Definition 10.1. *Consider ∂M endowed with a $\mathbf{CP}^1$ -structure, for instance coming from a complete hyperbolic metric on M . A **circle configuration** on ∂M is a finite set of open disks in ∂M (for the $\mathbf{CP}^1$ -structure), such that:*

- *the union of the closures of the disks is ∂M .*
- *the intersection points of the circles bounding two disks are never contained in any of the disks.*

Note that this definition implies that any point which is contained in two of the circles is in at least a third.

This relationship between ideal hyperbolic manifolds and circle configurations means that the results concerning the dihedral angles should be compared to those concerning configurations of circles, e.g. the rigidity results of [He99].

One can also consider circle packings in the more restrictive sense of sets of disks whose interiors are pairwise disjoint, with the complement made of disjoint, polygonal regions. Translating theorem 8.15, we will find the following result, which reduces to the classical Koebe circle packing theorem as extended by Thurston (see [Koe36, Thu97]).

Theorem 10.2. *Let Γ be the 1-skeleton of a triangulation of ∂M . There is a unique couple (g, c) , where g is a complete, convex co-compact hyperbolic metric on M , and c is a circle packing on ∂M (for the $\mathbf{CP}^1$ -structure defined on ∂M by g) whose incidence graph is Γ .*

=== Vérifier ref Thurston et al.

It is helpful to use a trick due to Thurston [Thu97]. Consider such a circle packing, such that the complement of the circles is the disjoint union of "triangles" bounded by 3 circle arcs. Then add, for each connected component of the complement of the circles, a circle which is orthogonal to the 3 circles which it interests. One then obtains a configuration of circles intersecting at right angles. The same construction works in the other way. Consider a configuration of circles intersecting at right angles, such that:

- the circles can be separated in two sets, the "white" and the "black" circles, such that each "black" circle only intersects transversally "white" circles, and conversely.
- each "black" circle intersects transversally exactly three "white" circles.

Then the set of "white" circles make up a circle packing.

Therefore, the proof theorem 10.2 is reduced to finding circle configurations on ∂M , where the intersections between the circles are always at right angles. Moreover, for the circle configuration coming from circle packings by adding orthogonal circles, it is easy to check that all circuits are made of at least 4 edges, and strictly more than 4 unless they are elementary. Therefore, theorem 8.15 applies, and theorem 10.2 follows.

If one considers graphs which are the 1-skeleton of cellulations (more general than triangulation) the same construction works; one obtains circles packings with the added property that, for each interstitial region, there is a circle orthogonal to all the adjacent circles. The uniqueness, under this additional condition, is proved by noting that each such packing gives rise to a circle configuration with right angle intersections, and thus to an ideal hyperbolic manifold with boundary faces intersecting at right angle (whose uniqueness is known by theorem 8.15).

Note that, seen in this light, the statements made here are related to different results on the rigidity of circle packings, see e.g. [Sch91].

In the fuchsian case, the results above can be stated as describing circle packings — or configurations of circles with given angles — on surfaces of genus $g \geq 2$.

10.2 Circles packings and CP^1 -structures

A slightly different idea is to consider the same circle packings on the boundary at infinity of $E(M)$, which, as a Riemann surface, can be endowed with a hyperbolic metric. The quasi-fuchsian case of our statements concerning dihedral angles should then be compared to recent results of S. Kojima, S. Mizushima and S. P. Tan [KMT01].

They consider a compact surface S of genus $g \geq 2$, with a triangulation τ . By a result of Thurston [Thu97], there is then a unique hyperbolic metric g_0 on S and a unique circle packing C_0 in g_0 whose nerve is τ ; it is called the Andreev-Thurston packing. Note that it can be recovered by applying the "Thurston trick" described above and theorem 8.15 in the fuchsian case.

Kojima, Mizushima and Tan are interested in the deformations of the CP^1 structure on S defined by g_0 , and of the circle packing C_0 . They prove among other things that, near (g_0, C_0) , there is a $6g - 6$ -parameter deformation space.

Notice that this local deformation result — which is only the easier part of the results of Kojima, Mizushima and Tan — can be recovered from the results of section 8, applied to the quasi-fuchsian case. Indeed, the Thurston-Andreev circle packing corresponds to a fuchsian ideal manifold M with all dihedral angles equal to $\pi/2$. Let Σ_+ and Σ_- be the two connected components of ∂M . Consider the deformations of the dihedral angles assignments on the edges of ∂M which do not change the angles on Σ_+ . This deformation space has dimension equal to the number of edges e of the triangulation τ , minus the number v of its vertices — because the sum at each vertex of the exterior dihedral angles has to remain 2π . But the Euler formula shows that $e - v = 6g - 6$.

Now each such deformation of the dihedral angles determines a unique quasi-fuchsian deformation of the hyperbolic structure on M by lemma 7.14; this in turn determines a unique deformation of the CP^1 structures on Σ_+ and Σ_- . Since the dihedral angles remain equal to $\pi/2$ on Σ_+ , the converse of the Thurston construction above still determines a circle packing, which is a deformation of C_0 . This recovers the deformation studied by Kojima, Mizushima and Tan.

11 Concluding remarks and questions

11.1 Manifolds with cusps

Most of what was done in this paper could presumably be extended from convex co-compact manifolds to manifolds with cusps; that is, we consider manifolds M which remain of finite volume, with a boundary which is polyhedral, i.e. the union of a finite set of totally geodesic faces which intersect along edges, with all their vertices ideal; but we allow those manifolds to have some cusps, i.e. M has finite volume but contains points arbitrarily far from the boundary.

The most delicate technical point to check is the content of section 4, concerning ideal triangulation. Proposition 5.5 extends to the case with cusps using the ideas of Epstein and Penner [EP88] in a manner which is closer to what they do in their paper, i.e. using the properties of the action of $\pi_1 M$ on the parabolic points. The combinatorial arguments used in section 4 to obtain an ideal triangulation from a cellulation then work without any difference in the case with cusps.

Doing this in the special case of a manifold with one cusp, one could presumably recover some results of Thurston on circle packings on the torus (see [Thu97], section 13.7). The dihedral angles, however, should not be restricted to be acute in this approach.

11.2 Hyperideal polyhedra

A remarkable point is that some of the properties described in the introduction survive when one considers polyhedra or surfaces with complete metrics of infinite area. In the polyhedral case this corresponds to hyperideal polyhedra; in the projective model of H^3 , they can be described as polyhedra having some "usual" vertices in H^3 and some "hyperideal" vertices beyond infinity, but such that all edges meet the interior of H^3 . An existence and uniqueness statement for the induced metric and the third fundamental form on those polyhedra can be found in [Sch98a], while a recent work of Bao and Bonahon [BB02] describes the set of their dihedral angles — a problem which is related to the third fundamental form.

For smooth surfaces, it looks like a possible analog of hyperideal polyhedra is the class of convex surfaces in H^3 whose boundary at infinity is a circle. A statement concerning the induced metrics and the third fundamental forms of such surfaces can be found in [Sch98b], but it deals only with the special case of surfaces of constant Gauss curvature. I also believe that one could prove a similar existence result for hyperbolic metrics on a manifold M inducing a given metric, with constant curvature $K \in [-1, 0)$, on the boundary (see [MS02]). A more general result, extending to metrics of non-constant curvature on the boundary, would be more demanding in terms of analytical techniques.

It would be interesting to understand whether those property also remain valid for hyperbolic manifolds with boundaries, when the metric has a "boundary at finite distance" and also a "boundary at infinity", thus extending the notion of hyperideal polyhedron. One might consider either a "polyhedral" type "finite boundary" — which would have to be "hyperideal" in the sense that, for each end, all edges "converge" to a hyperideal vertex — or a smooth "finite boundary", maybe with the additional condition that it meets the boundary at infinity along a circle.

One basic problem here is that we still lack an infinitesimal rigidity result here, stating that each infinitesimal deformation of those objects induces a non-trivial variation of both the induced metric and the third fundamental form. I believe that this could be achieved by using the same line of reasoning as here, but in the hyperideal instead of the ideal case. More precisely, the argument given at the end of section 2 to prove that the volume of ideal simplices is concave, also works for hyperideal simplices, if the volume of a hyperideal simplex is defined as the volume of the corresponding truncated simplex. Once this key point is obtained, the same method as here could presumably be used to understand "manifolds with hyperideal-like boundary". A rigidity result for hyperideal polyhedra, or for manifolds with boundaries locally like hyperideal polyhedra, would follow. I'm hoping to come back to this matter in the future.

11.3 Complete manifolds of finite volume

Some of the tools used here could also have applications to the studying the deformations of complete hyperbolic manifolds of finite volume.

Those manifold have a decomposition in cells, each isometric to an ideal polyhedron, by [EP88]. Thus the methods of section 5 might lead to ideal triangulations of finite covers, as in e.g. [NZ85].

One could then study the deformations obtained by changing the dihedral angles of the ideal triangles; the volume remains a concave functional on the space of angle assignments.

An important difference with the case of ideal hyperbolic manifolds, however, is that the shears along the singular edges is not the only obstruction to having a complete structure. The link of the vertices is now a torus — instead of a disk for ideal hyperbolic manifolds — which carries a similarity structure, with singularities at points corresponding to the singular edges; to have a complete structure, it is also necessary that the conformal part of the holonomy of each link vanishes. But this vanishing could also be a necessary and sufficient condition for the vanishing of the differential of the volume among deformations which do not change the total angle around each singular edge (at least it is sufficient).

11.4 Convex cores

The questions mentioned in the introduction, concerning the extension of theorems 1.1 and 1.2 to cases where ∂M is not smooth, have another natural setting: the case where M is supposed to be the convex core of a complete, convex co-compact manifolds N . This is equivalent to supposing that ∂M is convex and developpable, or that it is convex and has a hyperbolic (i.e. constant curvature -1) induced metric.

The question concerning theorem 1.1 is whether, for any hyperbolic metric h on ∂M , there is a unique hyperbolic metric g on M such that ∂M is convex and developpable, with induced metric h . The existence part holds, as was proved by Labourie [Lab91] (another proof exists, using a degree argument and the known results from [EM86] on a conjecture of Sullivan). The uniqueness, however, remains unknown.

When one considers the third fundamental forms of the boundary, one encounters a statement on the pleating measures of convex cores. Here again, satisfactory results on the existence part have been obtained recently by Bonahon, Otal [BO01] and Lecuire [Lec02], but the uniqueness is unknown.

Note that the questions on the convex cores can rather naturally be extended to the "horned hyperbolic manifolds" introduced in section 3. For instance, the question on the induced metric is:

Question 11.1. *Let h be a complete, finite area metric on ∂M minus a finite number of points. Is there a unique metric g on M such that (M, g) is a hyperbolic manifold with horns and that the induced metric on ∂M is h ?*

11.5 Orbifolds, cone-manifolds

The results given here, in particular theorem 8.15, can might also be used to construct hyperbolic orbifolds, or hyperbolic manifolds. This already happens in the fuchsian case, see [Thu97].

Other relation with cone-manifolds can be obtained, given an ideal hyperbolic manifold M , by gluing two copies of it along their common boundary. One obtains a finite volume, non-compact cone-manifold which is singular along a family of geodesics — corresponding to the edges of M — each going between two points at infinity. Moreover, the convexity of M means that the total angle around the singular geodesics is always strictly less than 2π . The infinitesimal rigidity of ideal hyperbolic manifolds could therefore be a consequence of an infinitesimal rigidity result for such cone-manifolds, which could perhaps be proved using the ideas of [HK98]. Moreover, those non-compact cone-manifolds could themselves be of interest.

11.6 Affine structures

There are several questions related to the affine structures coming from the dihedral angles of the boundary. It would be interesting to understand if the affine structure extends also to bent hyperbolic manifolds, so as to define an affine structure on the space of hyperbolic manifolds with horns.

Another question is whether the space of ideal hyperbolic manifolds — or even of hyperbolic manifolds with horns — is convex for this affine structure. A related question concerns the behavior near the boundary of the cellulation by the combinatorics of the boundary. For instance, is this cellulation locally finite at a boundary point corresponding to no bending and no degeneration of the conformal structure ?

11.7 Smooth boundaries

The questions treated in this paper have analogs for hyperbolic manifolds with smooth boundary. For instance, given a complete metric h of finite area, with $K > -1$ on ∂M minus a given number of points, which is asymptotically hyperbolic, is there a unique hyperbolic metric on M with convex boundary, inducing h on the boundary ?

The results quoted above on hyperideal manifolds might induce one to believe that the same question can be asked for metrics of infinite area, under the condition that the boundary at infinity of each end is a circle in $\partial_\infty M$. This is the analog of polyhedral condition that appeared in the previous sub-section. Even when M is a ball, however, this is proved only for constant curvature metrics on ∂M minus N points (see [Sch98b], which also contains a dual result on the third fundamental forms).

11.8 Higher dimensions

There is some hope of finding some properties related to questions A and B in higher dimension, in the setting of Einstein manifolds of negative curvature. An elementary step in this direction is taken in [Sch01b], where an infinitesimal deformation result concerning question A is proved.

Another similar perspective is given by the work of Graham and Lee [GL91], which can be considered as a first step towards a possible extension of the classical Ahlfors-Bers theorem in higher dimension, again in the setting of Einstein manifolds. Similar ideas form part of the mathematical side of the "AdS-CFT correspondence", an important conjecture in mathematical physics relating deep properties of conformally compact Einstein manifolds to other deep properties of their conformal boundaries; see e.g. [Mal98, Wit98, WY99, GW] and the references given there.

Acknowledgments

This work owes much to remarks made, at various times, by Francis Bonahon, François Labourie, Greg McShane and Igor Rivin. I also got a significant help from Richard Kenyon. Other helpful comments came from Michel Boileau, Philippe Eyssidieux, Gilbert Levitt, Anne Parreau and Joan Porti. I would like to thank them all here.

References

- [Ahl66] L. V. Ahlfors. *Lectures on quasiconformal mappings*. D. Van Nostrand Co., Inc., Toronto, Ont.-New York-London, 1966. Manuscript prepared with the assistance of Clifford J. Earle, Jr. Van Nostrand Mathematical Studies, No. 10.
- [Ale58] A. D. Aleksandrov. *Vestnik Leningrad Univ.*, 13(1), 1958.
- [And70] E.M. Andreev. Convex polyhedra in Lobacevskii space. *Mat. Sb.(N.S.)*, 81 (123):445–478, 1970.
- [And71] E.M. Andreev. On convex polyhedra of finite volume in Lobacevskii space. *Math. USSR Sbornik*, 12 (3):225–259, 1971.
- [BB02] X. Bao and F. Bonahon. Hyperideal polyhedra in hyperbolic 3-space. Preprint available at <http://math.usc.edu/~fbonahon>. *Bull. Soc. Math. France*, to appear, 2002.
- [Ber] Andreas Bernig. Variation of curvatures of subanalytic spaces and Schläfli-type formulas. Preprint 17-2002, U. Zürich.
- [BO01] F. Bonahon and J.-P. Otal. Laminations mesurées de plissage des variétés hyperboliques de dimension 3. <http://math.usc.edu/~fbonahon>, 2001.
- [Bon98a] F. Bonahon. A Schläfli-type formula for convex cores of hyperbolic 3-manifolds. *J. Differential Geom.*, 50(1):25–58, 1998.
- [Bon98b] F. Bonahon. Variations of the boundary geometry of 3-dimensional hyperbolic convex cores. *J. Differential Geom.*, 50(1):1–24, 1998.
- [Brä92] Walter Brägger. Kreispackungen und Triangulierungen. *Enseign. Math. (2)*, 38(3-4):201–217, 1992.
- [Cau13] A. L. Cauchy. Sur les polygones et polyèdres, second mémoire. *Journal de l'Ecole Polytechnique*, 19:87–98, 1813.
- [CD95] R. Charney and M. Davis. The polar dual of a convex polyhedral set in hyperbolic space. *Michigan Math. J.*, 42:479–510, 1995.
- [CdV91] Yves Colin de Verdière. Un principe variationnel pour les empilements de cercles. *Invent. Math.*, 104(3):655–669, 1991.
- [DM99] M. W. Davis and G. Moussong. Notes on nonpositively curved polyhedra. In *Low dimensional topology (Eger, 1996/Budapest, 1998)*, pages 11–94. János Bolyai Math. Soc., Budapest, 1999.
- [EM86] D. B. A. Epstein and A. Marden. Convex hulls in hyperbolic spaces, a theorem of Sullivan, and measured pleated surfaces. In D. B. A. Epstein, editor, *Analytical and geometric aspects of hyperbolic space*, volume 111 of *L.M.S. Lecture Note Series*. Cambridge University Press, 1986.
- [EP88] D. B. A. Epstein and R. C. Penner. Euclidean decompositions of noncompact hyperbolic manifolds. *J. Differential Geom.*, 27(1):67–80, 1988.

- [FLP91] A. Fathi, F. Laudenbach, and V. Poenaru. *Travaux de Thurston sur les surfaces*. Société Mathématique de France, Paris, 1991. Séminaire Orsay, Reprint of *Travaux de Thurston sur les surfaces*, Soc. Math. France, Paris, 1979 [MR 82m:57003], Astérisque No. 66-67 (1991).
- [GL91] C. R. Graham and J. M. Lee. Einstein metrics with prescribed conformal infinity on the ball. *Adv. Math.*, 87:186–225, 1991.
- [GW] C. R. Graham and E. Witten. Conformal anomaly of submanifold observables in AdS/CFT correspondence. hep-th/9901021.
- [He99] Zheng-Xu He. Rigidity of infinite disk patterns. *Ann. of Math. (2)*, 149(1):1–33, 1999.
- [HK98] C. D. Hodgson and S. P. Kerckhoff. Rigidity of hyperbolic cone-manifolds and hyperbolic Dehn surgery. *J. Differential Geom.*, 48:1–60, 1998.
- [Isk00] I. Iskhakov. *On hyperbolic surface tessellations and equivariant spacelike convex polyhedral surfaces in Minkowski space*. PhD thesis, Ohio State University, 2000.
- [KMT01] S. Kojima, S. Mizushima, and S. P. Tan. Circle packings on surfaces with projective structures. Available at: <http://www.is.titech.ac.jp/~sadayosi/publ/index-e.html>, 2001.
- [Koe36] P. Koebe. Kontaktprobleme der konformen Abbildung. *Abh. Sächs. Akad. Wiss. Leipzig Math.-Natur. Kl.*, 88:141–164, 1936.
- [Lab91] François Labourie. Problème de Minkowski et surfaces à courbure constante dans les variétés hyperboliques. *Bull. Soc. Math. France*, 119(3):307–325, 1991.
- [Lab92] F. Labourie. Métriques prescrites sur le bord des variétés hyperboliques de dimension 3. *J. Differential Geom.*, 35:609–626, 1992.
- [Lec02] C Lecuire. Plissage des variétés hyperboliques de dimension 3. Preprint 301, UMPA, ENS Lyon; <http://www.umpa.ens-lyon.fr/UMPA/Prepublications/liste.html>, 2002.
- [Lei01a] G. Leibon. Characterizing the Delaunay decompositions of compact hyperbolic surfaces. math.GT/0103174, 2001.
- [Lei01b] G. Leibon. Random delaunay triangulations, the Thurston-Andreev theorem, and metric uniformization. math.DG/0011016, 2001.
- [LS00] F. Labourie and Jean-Marc Schlenker. Surfaces convexes fuchsienues dans les espaces lorentziens à courbure constante. *Math. Annalen*, 316:465–483, 2000.
- [LS01] Roger C. Lyndon and Paul E. Schupp. *Combinatorial group theory*. Springer-Verlag, Berlin, 2001. Reprint of the 1977 edition.
- [Mal98] J. Maldacena. The large N limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.*, 2:231–252, 1998.
- [Mil94] J. Milnor. The Schläfli differential equality. In *Collected papers, vol. 1*. Publish or Perish, 1994.
- [MS02] Greg McShane and Jean-Marc Schlenker. Hyperbolic manifolds with constant curvature boundaries. In preparation, 2002.
- [NZ85] Walter D. Neumann and Don Zagier. Volumes of hyperbolic three-manifolds. *Topology*, 24(3):307–332, 1985.
- [Pog73] A. V. Pogorelov. *Extrinsic Geometry of Convex Surfaces*. American Mathematical Society, 1973. Translations of Mathematical Monographs. Vol. 35.
- [PP00] Carlo Petronio and Joan Porti. Negatively oriented ideal triangulations and a proof of Thurston’s hyperbolic Dehn filling theorem. *Expo. Math.*, 18(1):1–35, 2000.

- [RH93] I. Rivin and C. D. Hodgson. A characterization of compact convex polyhedra in hyperbolic 3-space. *Invent. Math.*, 111:77–111, 1993.
- [Riv86] I. Rivin. *Thesis*. PhD thesis, Princeton University, 1986.
- [Riv92] I. Rivin. Intrinsic geometry of convex ideal polyhedra in hyperbolic 3-space. In M. Gyllenberg and L. E. Persson, editors, *Analysis, Algebra, and Computers in Mathematical Research*, pages 275–292. Marcel Dekker, 1992. (Proc. of the 21st Nordic Congress of Mathematicians).
- [Riv94] I. Rivin. Euclidean structures on simplicial surfaces and hyperbolic volume. *Annals of Math.*, 139:553–580, 1994.
- [Riv96] I. Rivin. A characterization of ideal polyhedra in hyperbolic 3-space. *Annals of Math.*, 143:51–70, 1996.
- [Rou02] Mathias Rousset. Polyèdres hyperidéaux fuchsien. Rapport de stage d’option de l’Ecole Polytechnique, promo X99, 2002.
- [RS99] Igor Rivin and Jean-Marc Schlenker. The Schläfli formula in Einstein manifolds with boundary. *Electronic Research Announcements of the A.M.S.*, 5:18–23, 1999.
- [RS00] Igor Rivin and Jean-Marc Schlenker. The Schläfli formula and Einstein manifolds. Preprint math.DG/0001176, 2000.
- [Sch91] Oded Schramm. Rigidity of infinite (circle) packings. *J. Amer. Math. Soc.*, 4(1):127–149, 1991.
- [Sch94] Jean-Marc Schlenker. Surfaces elliptiques dans des espaces lorentziens à courbure constante. *Compte Rendus de l’Académie des Sciences, Série A*, 319:609–614, 1994.
- [Sch96] Jean-Marc Schlenker. Surfaces convexes dans des espaces lorentziens à courbure constante. *Commun. Anal. and Geom.*, 4:285–331, 1996.
- [Sch98a] Jean-Marc Schlenker. Métriques sur les polyèdres hyperboliques convexes. *J. Differential Geom.*, 48(2):323–405, 1998.
- [Sch98b] Jean-Marc Schlenker. Représentations de surfaces hyperboliques complètes dans H^3 . *Annales de l’Institut Fourier*, 48(3):837–860, 1998.
- [Sch00] Jean-Marc Schlenker. Dihedral angles of convex polyhedra. *Discrete Comput. Geom.*, 23(3):409–417, 2000.
- [Sch01a] Jean-Marc Schlenker. Convex polyhedra in Lorentzian space-forms. *Asian J. of Math.*, 5:327–364, 2001.
- [Sch01b] Jean-Marc Schlenker. Einstein manifolds with convex boundaries. *Commentarii Math. Helvetici*, 76(1):1–28, 2001.
- [Sch02a] Jean-Marc Schlenker. Hyperbolic manifolds with convex boundary. preprint, math.DG/0205305, available at <http://picard.ups-tlse.fr/~schlenker>, 2002.
- [Sch02b] Jean-Marc Schlenker. Hypersurfaces in H^n and the space of its horospheres. *Geom. Funct. Anal.*, 12:395–435, 2002.
- [SP00] Eva Suárez-Peiró. A Schläfli differential formula for simplices in semi-Riemannian hyperquadrics, Gauss-Bonnet formulas for simplices in the de Sitter sphere and the dual volume of a hyperbolic simplex. *Pacific J. Math.*, 194(1):229–255, 2000.
- [Sto68] J. J. Stoker. Geometrical problems concerning polyhedra in the large. *Comm. Pure Appl. Math.*, 21:119–168, 1968.

- [Thu97] William P. Thurston. Three-dimensional geometry and topology. Available on the MSRI website, at: <http://www.msri.org/publications/books/gt3m/>; recent version of the 1980 notes, 1997.
- [Tro91] Marc Troyanov. Prescribing curvature on compact surfaces with conical singularities. *Trans. Amer. Math. Soc.*, 324(2):793–821, 1991.
- [Vin93] E.B. Vinberg, editor. *Geometry II, Geometry of Spaces of Constant Curvature*, volume 29 of *Encyclopaedia of Mathematical Sciences*. Springer, 1993.
- [Wit98] E. Witten. Anti de Sitter space and holography. *Adv. Theor. Math. Phys.*, 2:253–291, 1998.
- [WY99] Edward Witten and S.-T. Yau. Connectedness of the boundary in the AdS/CFT correspondence. *Adv. Theor. Math. Phys.*, 3(6):1635–1655 (2000), 1999. hep-th/9910245.